

BIRCH, STEWART, KOLASCH & BIRCH, LLP

INTELLECTUAL PROPERTY LAW

8110 GATEHOUSE ROAD

SUITE 500 EAST

FALLS CHURCH, VA 22042

USA

(703) 205-8000

FAX: (703) 205-8050

(703) 698-8590 (G IV)

e-mail: mailroom@bskb.com

web: http://www.bskb.com

GARY D. YACURA  
THOMAS S. AUCHT  
MICHAEL R. CAMMA  
JAMES T. ELLER, JR.  
SCOTT L. LOWE  
JOSEPH H. KIM, Ph.D.  
RICHARD S. MYERS, JR.\*  
MARY ANN CAPRIA  
MICHAEL J. CORNELISON\*  
MARK J. NUEL, Ph.D.  
ROBERT V. RACUNAS  
DARIN E. BARTHOLOMEW\*  
D. RICHARD ANDERSON  
PAUL C. LEWIS  
JERRY W. HOGGE

REG. PATENT AGENTS:  
FREDERICK R. HANDREN  
ANDREW J. TELESZ, JR.  
MARYANNE LIOTTA, Ph.D.  
MAKI HATSUMI  
STEVEN P. WIGMORE  
ESTHER H. CHIN  
MIKE S. RYU  
W. KARL RENNER  
CRAIG A. MCROBBIE  
GARTH M. DAHLEN, Ph.D.  
LAURA C. LUTZ  
ROBERT E. GOOZNER, Ph.D.  
HYUNG N. SOHN

TERRELL C. BIRCH  
RAYMOND C. STEWART  
JOSEPH A. KOLASCH  
JAMES M. SLATTERY  
BERNARD L. SWEENEY\*  
MICHAEL K. MUTTER  
CHARLES GORENSTEIN  
GERALD M. MURPHY, JR.  
LEONARD R. SVENSSON  
TERRY L. CLARK  
ANDREW D. MEIKLE  
MARC S. WEINER  
JOE MCKINNEY MUNCY  
ROBERT J. KENNEY  
JOSEPH FARACI  
DONALD J. DALEY  
JOHN W. BAILEY  
JOHN A. CASTELLANO, III

SENIOR COUNSEL:  
ANTHONY L. BIRCH

OF COUNSEL:  
HERBERT M. BIRCH  
(1905-1996)  
ELLIOT A. GOLDBERG\*  
WILLIAM L. GATES\*  
EDWARD H. VALANCE  
RUPERT J. BRADY (RET.)\*

\*ADMITTED TO A BAR OTHER THAN VA.

Date: April 29, 1999

Docket No.: 0020-4559P

Assistant Commissioner for Patents  
Box PATENT APPLICATION  
Washington, D.C. 20231

Sir:

Transmitted herewith for filing is the patent application of

Inventor(s): WATANABE, Eijiro  
OEDA, Kenji

For: RAFFINOSE SYNTHASE GENES AND THEIR USE

Enclosed are:

- X  A specification consisting of  100  pages  
\_\_\_\_ sheet(s) of \_\_\_\_\_ drawings  
 X  An assignment of the invention  
 X  Certified copy of Priority Document(s)  
 X  Executed Declaration  X  Original \_\_\_\_ Photocopy  
\_\_\_\_ A verified statement to establish small entity status under 37  
CFR 1.9 and 37 CFR 1.27  
 X  Preliminary Amendment  
\_\_\_\_ Information Disclosure Statement, PTO-1449 and reference(s)

X Other Substitute Sequence Listing with disk

The filing fee has been calculated as shown below:

LARGE ENTITY				SMALL ENTITY	
FOR	NO. FILED	NO. EXTRA	RATE FEE		RATE FEE
BASIC FEE	***** ***** *****	***** ***** *****	***** ***** \$760.00 *****	or	**** **** \$380.00 ****
TOTAL CLAIMS	27 - 20 =	7	x18 =\$ 126.00	or	x 9 = \$ 0.00
INDEPENDENT	14 - 3 =	11	x78 =\$ 858.00	or	x 39 = \$ 0.00
MULTIPLE DEPENDENT CLAIM PRESENTED <u>no</u>			+260 = \$ 0.00	or	+130 = \$ 0.00
			TOTAL \$1,744.00		TOTAL \$ 0.00

✓ X A check in the amount of \$1,784.00 to cover the filing fee and recording fee (if applicable) is enclosed.


— Please charge Deposit Account No. 02-2448 in the amount of \$\_\_\_\_\_. A triplicate copy of this transmittal form is enclosed.

— No fee is enclosed.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. 1.16 or under 37 C.F.R. 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By   
GERALD M. MURPHY, JR.,  
Reg. No. 28,977  
P. O. Box 747  
Falls Church, Virginia 22040-0747

**BOX SEQUENCE**  
**PATENT**  
**20-4559P**

IN THE U.S. PATENT AND TRADEMARK OFFICE

APPLICANT(S): Eijiro WATANABE et al.

APPLICATION NO.: New

GROUP: Not Assigned

FILED: April 29, 1999

EXAMINER: Not Assigned

FOR: RAFFINOSE SYNTHASE GENES AND THEIR USE

**PRELIMINARY AMENDMENT SUPPLEMENTAL TO SEQUENCE LISTING**

Honorable Commissioner of Patents  
Washington, D.C. 20231

April 29, 1999

Sir:

The following preliminary amendments and remarks are respectfully submitted in connection with the above-identified application.

***IN THE SPECIFICATION***

Please amend the specification as follows:

Page 10

Line 10, after "11" insert -- (SEQ ID NO:9)--.

Line 10, after "12" insert -- (SEQ ID NO:10)--.

Page 11

Line 9, after “11” insert -- (SEQ ID NO:9)--.

Line 10 after “12” insert -- (SEQ ID NO:10)--.

Page 12

Line 5, after “21” insert -- (SEQ ID NO:11)--.

Line 5, after “22” insert -- (SEQ ID NO:12)--.

Line 19, after “23” insert -- (SEQ ID NO:13)--.

Line 19, after “24” insert -- (SEQ ID NO:14)--.

Page 13

Line 13, after “21” insert -- (SEQ ID NO:11)--.

Line 14, after “22” insert -- (SEQ ID NO:12)--.

Line 15, after “23” insert -- (SEQ ID NO:13)--.

Line 16, after “24” insert -- (SEQ ID NO:14)--.

Page 14

Line 13, after “33” insert -- (SEQ ID NO:17)--.

Line 13, after “34” insert -- (SEQ ID NO:18)--.

Page 15

Line 5, after “35” insert -- (SEQ ID NO:19)--.

Line 5, after “36” insert -- (SEQ ID NO:20)--.

Line 23, after “31” insert -- (SEQ ID NO:15)--.

Line 24, after “32” insert -- (SEQ ID NO:16)--.

Line 25, after “33” insert -- (SEQ ID NO:17)--.

Page 16

Line 1, after “34” insert -- (SEQ ID NO:18)--.

Line 2, after “35” insert -- (SEQ ID NO:19)--.

Line 3, after “36” insert -- (SEQ ID NO:20)--.

Page 25

Line 4, after “31” insert -- (SEQ ID NO:15)--.

Line 5, after “32” insert -- (SEQ ID NO:16)--.

Page 39

Line 23, after “List 4” insert -- (SEQ ID NOS:21 & 22)--.

Page 40

Line 10, after “List 5” insert -- (SEQ ID NO:23)--.

Line 16, after “List 5” insert -- (SEQ ID NO:23)--.

Line 22, after “primer” insert -- (SEQ ID NO:21)--.

Line 24, after “primer” insert -- (SEQ ID NO:22)--.

Page 41

Line 3, after “5-SC-2” insert -- (SEQ ID NO:23)--.

Page 42

Line 11, after “List 6” insert -- (SEQ ID NOS:24-27)--.

Line 14, after “primers” insert -- (SEQ ID NOS:24-27)--.

Line 18, after “6-3-F” insert -- (SEQ ID NO:24)--.

Line 18, after “6-8-RV” insert -- (SEQ ID NO:25)--.

Line 18, after “6-10-F” insert -- (SEQ ID NO:26)--.

Line 18, after “6-6-RV” insert -- (SEQ ID NO:27)--.

Page 43

Line 1, after “List 7” insert -- (SEQ ID NOS:28-35)--.

Line 7, after “6-3-F” insert -- (SEQ ID NO:24)--.

Line 9, after “6-8-RV” insert -- (SEQ ID NO:25)--.

Line 11, after “6-10-F” insert -- (SEQ ID NO:26)--.

Line 13, after “6-6-RV” insert -- (SEQ ID NO:27)--.

Line 16, after “7-Sb-1” insert -- (SEQ ID NO:28)--.

Line 18, after “7-Sb-2RV” insert -- (SEQ ID NO:29)--.

Line 20, after “7-Sb-3RV” insert -- (SEQ ID NO:30)--.

Line 22, after “7-Sb-4RV” insert -- (SEQ ID NO:31)--.

Line 24, after “7-Sb-5” insert -- (SEQ ID NO:32)--.

Page 44

Line 1, after “7-Sb-6” insert -- (SEQ ID NO:33)--.

Line 3, after “7-Sb-7” insert -- (SEQ ID NO:34)--.

Line 5, after “7-Sb-8RV” insert -- (SEQ ID NO:35)--.

Page 45

Line 16, after “List 8” insert -- (SEQ ID NOS:36 & 37)--.

Line 19, after “primers” insert -- (SEQ ID NOS:36 & 37)--.

Page 46

Line 5, after “List 9” insert -- (SEQ ID NOS:38 & 39)--.

Line 14, after “8-#1” insert -- (SEQ ID NO:36)--.

Line 16, after “8-#10RV” insert -- (SEQ ID NO:37)--.

Line 19, after “9-primer-1” insert -- (SEQ ID NO:38)--.

Line 21, after “9-primer-2RV” insert -- (SEQ ID NO:39)--.

Page 47

Line 3, after “6” insert -- (SEQ ID NOS:6, 8, 36-39)--.

Line 4, after “10” insert -- (SEQ ID NOS:40-46)--.

Line 9, after “10-B-2RV” insert -- (SEQ ID NO:40)--.

Line 9, after “10-B-3RV” insert -- (SEQ ID NO:41)--.

Line 9, after “10-B-4RV” insert -- (SEQ ID NO:42)--.

Line 10, after “10-B-1” insert -- (SEQ ID NO:43)--.

Line 10, after “10-B-8” insert -- (SEQ ID NO:44)--.

Line 10, after “10-B-7” insert -- (SEQ ID NO:45)--.

Line 11, after “10-B-6” insert -- (SEQ ID NO:46)--.

Line 18, after “10-B-2RV” insert -- (SEQ ID NO:40)--.

Line 20, after “10-B-3RV” insert -- (SEQ ID NO:41)--.

Line 22, after “10-B-4RV” insert -- (SEQ ID NO:42)--.

Line 24, after “10-B-1” insert -- (SEQ ID NO:43)--.

Page 48

Line 2, after “10-B-8” insert -- (SEQ ID NO:44)--.

Line 4, after “10-B-7” insert -- (SEQ ID NO:45)--.

Line 6, after “10-B-6” insert -- (SEQ ID NO:46)--.

Line 13, after “List 11” insert -- (SEQ ID NOS:47 & 48)--.

Line 19, after “List 12” insert -- (SEQ ID NOS:49 & 50)--.

Line 22, after “List 13” insert -- (SEQ ID NOS:51-53)--.

Page 49

Line 4, after “11-SacI-BjN” insert -- (SEQ ID NO:47)--.

Line 6, after “11-SacI-BjintRV” insert -- (SEQ ID NO:48)--.

Line 9, after “12-BamSac-(+)linker” insert -- (SEQ ID NO:49)--.

Line 11, after “12-BamSac-(-)linker” insert -- (SEQ ID NO:50)--.

Line 14, after “13-35S-3” insert -- (SEQ ID NO:51)--.



Line 16, after "13-B-2RV" insert -- (SEQ ID NO:52)--.

Line 18, after "13-B-8" insert -- (SEQ ID NO:53)--.

Page 52

Line 19, after "List 1" insert -- (SEQ ID NOS:9 & 10)--.

Line 22, after "11" insert -- (SEQ ID NO:9)--.

Line 23, after "12" insert -- (SEQ ID NO:10)--.

Page 53

Line 2, after "List 2" insert -- (SEQ ID NOS:11-14)--.

Line 4, after "21" insert -- (SEQ ID NO:11)--.

Line 6, after "22" insert -- (SEQ ID NO:12)--.

Line 10, after "23" insert -- (SEQ ID NO:13)--.

Line 11, after "24" insert -- (SEQ ID NO:14)--.

Line 15, after "31," insert -- (SEQ ID NO:15)--.

Line 15, after "32" insert -- (SEQ ID NO:16)--.

Line 17, after "31" insert -- (SEQ ID NO:15)--.

Line 19, after "32" insert -- (SEQ ID NO:16)--.

Line 23, after "33" insert -- (SEQ ID NO:17)--.

Line 23, after "34" insert -- (SEQ ID NO:18)--.

Line 25, after "33" insert -- (SEQ ID NO:17)--.

Line 25, after "34" insert -- (SEQ ID NO:18)--.

Page 54

Line 1, after “33” insert -- (SEQ ID NO:17)--.

Line 3, after “34” insert -- (SEQ ID NO:18)--.

Line 6, after “35” insert -- (SEQ ID NO:19)--.

Line 6, after “36” insert -- (SEQ ID NO:20)--.

Line 9, after “35” insert -- (SEQ ID NO:19)--.

Line 10, after “36” insert -- (SEQ ID NO:20)--.

Line 16, after “List 4” insert -- (SEQ ID NOS:21 & 22)--.

Line 24, after “List 5” insert -- (SEQ ID NO:23)--.

Page 55

Line 1, after “5-SC-2” insert -- (SEQ ID NO:23)--.

Line 4, after “List 6” insert -- (SEQ ID NOS:24-27)--.

Line 12, after “List 7” insert -- (SEQ ID NOS:28-35)--.

Line 18, after “List 8” insert -- (SEQ ID NOS:36 & 37)--.

Page 56

Line 1, after “List 9” insert -- (SEQ ID NOS:38 & 39)--.

Line 7, after “List 10” insert -- (SEQ ID NOS:40-46)--.

Line 13, after “List 11” insert -- (SEQ ID NOS:47 & 48)--.

Line 15, after “11-SacI-BjN” insert -- (SEQ ID NO:47)--.

Line 18, after “11-SacI-BjintRV” insert -- (SEQ ID NO:48)--.

Line 22, after “List 12” insert -- (SEQ ID NOS:49 & 50)--.

Page 57

Line 5, after "List 13" insert -- (SEQ ID NOS:51-53)--.

Line 7, after "13-35S-3" insert -- (SEQ ID NO:51)--.

Line 8, after "13-B-2RV" insert -- (SEQ ID NO:52)--.

Line 10, after "13-B-8" insert -- (SEQ ID NO:53)--.

Please delete pages 59 through 94 of the specification containing the Sequence Listing.  
Please renumber the remaining pages of the specification, beginning with the claims,  
consecutively from page 59 of the specification. Please insert the Substitute Sequence Listing  
enclosed herewith immediately after the abstract.

IN THE CLAIMS:

Claim 11, lines 2 and 3, change "any one of claims 1 to 10" to --claim 1--

Claim 16, lines 2 and 3, change "any one of claims 1 to 10" to --claim 1--

Claim 17, line 2, change "any one of claims 1 to 10" to --claim 1--

Claim 18, line 2, change "any one of claims 1 to 10" to --claim 1--

Claim 21, line 1, change "any one of claims 18 to 20" to --claim 18--

Claim 22, line 1, change "any one of claims 18 to 20" to --claim 18--

Claim 23, lines 3 and 4, change "any one of claims 18 to 22" to --18--


**REMARKS**

Enclosed herewith in full compliance to 37 C.F.R. §§1.821-1.825 is a Substitute Sequence Listing to be inserted into the specification as indicated above. The Substitute Sequence Listing in no way introduces new matter into the specification. Also submitted herewith in full compliance to 37 C.F.R. §§1.821-1.825 is a disk copy of the Substitute Sequence Listing. The disk copy of the Substitute Sequence Listing, file "20-4559P.txt", is identical to the paper copy, except that it lacks formatting. The amendment to the claims corrects any improper multiple dependent claims to place the application into better form.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future submissions, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. §1.16 or under 37 C.F.R. §1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By   
GERALD M. MURPHY, JR.  
Reg. No. 28,977  
P.O. Box 747  
Falls Church, VA 22040-0747  
(703) 205-8000

GMM/LPS

20-4559P

Enclosures: Paper Copy of Substitute Sequence Listing

Disk Copy of Substitute Sequence listing

## RAFFINOSE SYNTHASE GENES AND THEIR USE

### BACKGROUND OF THE INVENTION

#### 5 FIELD OF INVENTION

The present invention relates to raffinose synthase genes and their use.

#### DISCLOSURE OF THE RELATED ART

10 Raffinose family oligosaccharides are derivatives of sucrose, which are represented by the general formula:  $\alpha$ -D-galactopyranosyl-(1 $\rightarrow$ 6)<sub>n</sub>- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-fluctofuranoside, and they are called "raffinose" when n is 1, "stachyose" when n is 2, "verbascose" when n is 3, and "ajugose" when n is 4.

15 It has been known that raffinose family oligosaccharides have an effect of giving good conditions of enterobacterial flora, if present at an appropriate amount in food. Therefore, raffinose family oligosaccharides have already been used as a functional food material for addition  
20 to some kinds of food and utilized in the field of specific health food. On the other hand, raffinose family oligosaccharides are neither digested nor absorbed in mammals such as human, but are assimilated and decomposed by enterobacteria to generate gases and to cause meteorism and absorption disorder. Therefore, it  
25 has been desired to appropriately regulate the amount of raffinose family oligosaccharides in food and feed.

Raffinose family oligosaccharides are synthesized by the raffinose family oligosaccharide biosynthesis system beginning with sucrose in many plants. This biosynthesis system normally involves a reaction for the sequential addition of galactosyl groups from galactinol through an  $\alpha$  (1 $\rightarrow$ 6) bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule. Raffinose synthase is the enzyme concerned in the reaction for producing raffinose by allowing a D-galactosyl group derived from galactinol to form the  $\alpha$  (1 $\rightarrow$ 6) bond with the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule in the first step of this biosynthesis system. It has been suggested that this enzyme constitutes a rate-limiting step in the above synthesis system, and therefore this enzyme is quite important in the control of biosynthesis of raffinose family oligosaccharides.

Then, a method for controlling an expression level or activity of raffinose synthase in plants by utilizing a raffinose synthase gene is effective to control a biosynthesis system of raffinose family oligosaccharides in plants to increase or decrease the production of raffinose in plants.

Thus, a raffinose synthase gene which can be used in such a method has been desired.

The main object of the present invention is to provide novel raffinose synthase genes from plants.

This object as well as other objects and advantage of the present invention will become apparent to those skilled  
5 in the art from the following description.

#### SUMMARY OF THE INVENTION

Under these circumstances, the present inventors have intensively studied and succeeded in isolating novel genes  
10 encoding raffinose synthase from various plants. Thus, the present invention has been completed.

That is, the present invention provides:

1. A raffinose synthase gene which comprises a nucleotide sequence hybridizable with a nucleotide sequence  
15 selected from the group consisting of:

(a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1,

(b) the nucleotide sequence represented by SEQ ID NO: 2,

20 (c) a nucleotide sequence encoding the amino acid sequence of represented by SEQ ID NO: 3,

(d) the nucleotide sequence represented by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4,

25 (e) a nucleotide sequence encoding the amino acid

sequence represented by SEQ ID NO: 5,

(f) the nucleotide sequence represented by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6,

5 (g) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7, and

(h) the nucleotide sequence represented by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8,

10 under stringent conditions, and encoding a protein being capable of binding D-galactosyl group through  $\alpha$  (1 $\rightarrow$ 6) bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule to form raffinose.

2. A raffinose synthase gene comprising a  
15 nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1.

3. A raffinose synthase gene comprising the nucleotide sequence represented by SEQ ID NO: 2.

4. A raffinose synthase gene comprising a  
20 nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 3.

5. A raffinose synthase gene comprising the nucleotide sequence represented by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID  
25 NO: 4.



6. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5.

5 7. A raffinose synthase gene comprising the nucleotide sequence represented by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6.

10 8. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7.

9. A raffinose synthase gene comprising the nucleotide sequence represented by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8.

15 10. A raffinose synthase gene comprising the nucleotide sequence represented by SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8.

11. A nucleic acid comprising a partial nucleotide sequence of the raffinose synthase gene of any one of the above 1 to 10.

20 12. A method for detecting a nucleic acid containing a raffinose synthase gene which comprises detecting said nucleic acid by hybridization using the labeled nucleic acid of the above 11 as a probe.

25 13. A method for amplifying a nucleic acid containing a raffinose synthase gene which comprises amplifying

said nucleic acid by polymerase chain reaction (PCR) using the nucleic acid of the above 11 as a primer.

14. A method for obtaining a raffinose synthase gene which comprises the steps of:

5 detecting a nucleic acid containing said raffinose synthase gene by hybridization using the labeled nucleic acid of the above 11 as a probe, and

recovering the detected nucleic acid.

10 15. A method for obtaining a raffinose synthase gene which comprises the steps of:

amplifying a nucleic acid containing said raffinose synthase gene by PCR using the nucleic acid of the above 11 as a primer, and

recovering the amplified nucleic acid.

15 16. A nucleic acid comprising a nucleic acid containing the raffinose synthase gene of any one of the above 1 to 10 which is joined to a nucleic acid exhibiting promoter activity in a host cell.

20 17. A vector comprising the raffinose synthase gene of any one of the above 1 to 10.

18. A transformant, wherein the raffinose synthase gene of any one of the above 1 to 10 is introduced into a host cell.

25 19. A transformant, wherein the nucleic acid of the above 16 is introduced into a host cell.

20. A transformant, wherein the vector of the above 17 is introduced into a host cell.

21. The transformant of any one of the above 18 to 20, wherein the host is a microorganism.

5 22. The transformant of any one of the above 18 to 20, wherein the host is a plant.

23. A method for producing a raffinose synthase which comprises the steps of:

10 culturing or growing the transformant of any one of the above 18 to 22 to produce the raffinose synthase, and collecting the raffinose synthase.

24. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 1.

15 25. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 3.

26. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 5.

27. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 7.

20 The term "nucleic acid" used herein means an oligomer compound or a high molecular compound which is generally called "DNA" or "RNA".

#### DETAILED DESCRIPTION OF THE INVENTION

25 The gene engineering techniques described below can

be carried out, for example, according to methods described in  
"Molecular Cloning: A Laboratory Manual 2nd edition" (1989),  
Cold Spring Harbor Laboratory Press, ISBN 0-87969-309-6;

"Current Protocols In Molecular Biology" (1987), John Wiley &  
5 Sons, Inc. ISBN 0-471-50338-X; "Current Protocols In Protein  
Science" (1995), John Wiley & Sons, Inc. ISBN 0-471-11184-8.

The genes of the present invention can be obtained  
from soybean, plants belonging to the families Chenopdiaceae  
such as beet, etc. and Cruciferae such as mustard, rapeseed,  
10 etc. Specific examples of the genes of the present invention  
include those comprising a nucleotide sequence encoding the  
amino acid sequence represented by SEQ ID NO: 1, the nucleotide  
sequence represented by SEQ ID NO: 2, a nucleotide sequence  
encoding the amino acid sequence represented by SEQ ID NO: 3,  
15 the nucleotide sequence represented by SEQ ID NO: 4 or by the  
236th to 2584th nucleotides in the nucleotide sequence  
represented by SEQ ID NO: 4, a nucleotide sequence encoding the  
amino acid sequence represented by SEQ ID NO: 5, the nucleotide  
sequence represented by SEQ ID NO: 6 or by the 134th to 2467th  
20 nucleotides in the nucleotide sequence represented by SEQ ID  
NO: 6, a nucleotide sequence encoding the amino acid sequence  
represented by SEQ ID NO: 7, the nucleotide sequence represented  
by SEQ ID NO: 8 or by the 1st to 1719th nucleotides in the  
nucleotide sequence represented by SEQ ID NO: 8, and the like.

25 The genes of the present invention can be obtained,

for example, by the following method.

That is, the genes of the present invention derived from soybean can be obtained, for example, by the following method.

5           For example, the gene can be obtained by a hybridization method using a nucleic acid having the nucleotide sequence represented by SEQ ID NO: 2 or its partial nucleotide sequence as a probe to detect a nucleic acid fragment which hybridizes to the probe in DNAs derived from soybean, followed  
10 by isolating the detected nucleic acid.

          In this method, first, a nucleic acid to be used as the probe is prepared. As such a nucleic acid, for example, there is a nucleic acid composed of an oligonucleotide chemically synthesized by a conventional method on the basis of the  
15 nucleotide sequence of SEQ ID NO: 2. Specific example thereof includes a nucleic acid having the 800th to the 899th nucleotides in the nucleotide sequence represented by SEQ ID NO: 2.

          Alternatively, the gene of the present invention derived from soybean can be obtained by the following method.

20           For example, tissue of soybean (*Glycine max*) is frozen in liquid nitrogen and ground physically with a mortar or other means into finely divided tissue debris powder. From the tissue debris powder, RNA is extracted by a conventional method. A commercially available RNA extraction kit can be utilized in  
25 the extraction. RNA is recovered from thus-obtained RNA extract

by ethanol precipitation. Poly-A tailed RNA is fractionated from thus-recovered RNA by a conventional method. A commercially available oligo-dT column can be utilized in this fractionation. cDNA is synthesized from the poly-A tailed RNA thus obtained by a conventional method. The synthesis can be carried out by using a commercially available cDNA synthesis kit. DNA is amplified by PCR using the above-obtained cDNA as the template and primers designed and synthesized on the basis of the nucleotide sequence of SEQ ID NO: 2. More specifically, as the primers, for example, there are primers 11 and 12 shown in List 1 hereinafter. When PCR is carried out by using these primers and as the template cDNA derived from soybean, the genes of the present invention derived from soybean, e.g., the "raffinose synthase gene having a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 1" and the "raffinose synthase gene having the nucleotide sequence of SEQ ID No: 2" can be obtained.

The amplified DNA can be cloned according to a conventional method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; or "Current Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X.

Alternatively, cloning can be carried out, for example, by using a commercially available cloning kit such as TA cloning kit (Invitrogen) and a commercially available plasmid vector such

as pBluescript II (Stratagene). The nucleotide sequence of the DNA clone can be determined by dideoxy terminating method such as that described by F. Sanger, S. Nicklen, A.R. Coulson, Proceedings of National Academy of Science U.S.A. (1977), 74, pp. 5463-5467. For example, preferably, a commercially available kit such as ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit manufactured by Perkin-Elmer can be used.

List 1

Primer 11: ccaatctgat catgcttgtg ccgaa 25mer

10 Primer 12: ggaacaaagt tatgcactat tattttaaggt 30mer

The genes of the present invention derived from a Chenopdiaceae plant such as beet can be obtained by the following method.

For example, tissue of a Chenopdiaceae plant such as beet (*Beta vulgaris*) is frozen in liquid nitrogen and ground physically with a mortar or other means into finely divided tissue debris powder. From this tissue debris powder, RNA is extracted by a conventional method. A commercially available RNA extraction kit can be utilized in the extraction. RNA is recovered from the thus-obtained RNA extract by ethanol precipitation. From the recovered RNA, poly-A tailed RNA is fractionated by a conventional method. A commercially available oligo-dT column can be utilized in this fractionation.

cDNA is synthesized from the poly-A tailed RNA thus obtained by a conventional method. The synthesis can be carried out by

utilizing a commercially available cDNA synthesis kit. DNA is amplified by PCR using the above-obtained cDNA as the template and primers designed and chemically synthesized on the basis of the nucleotide sequence of SEQ ID NO: 4. More specifically, as the primers, for example, there are primers 21 and 22 shown in List 2 hereinafter. When PCR is carried out by using these primers and as the template cDNA derived from beet, the genes of the present invention derived from beet, e.g., the "raffinose synthase gene having a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 3," and the "raffinose synthase gene having a nucleotide sequence of SEQ ID No: 4" can be obtained.

According to a particular purpose, the PCR primers can also be designed and synthesized on the basis of the nucleotide sequence of SEQ ID NO: 4. For example, in order to amplify the "raffinose synthase gene having the nucleotide sequence represented by the 236th to the 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4", preferably, oligonucleotides having the nucleotide sequences represented by primers 23 and 24 in List 2 below are synthesized and used as the primers.

The amplified DNA can be cloned according to a conventional method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; or "Current Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X.



Alternatively, cloning can be carried out by using a commercially available cloning kit such as TA cloning kit (Invitrogen) and a commercially available plasmid vector such as pBluescript II (Stratagene). The nucleotide sequence of the DNA clone can be determined, for example, by dideoxy terminating method such as that described by F. Sanger, S. Nicklen, A.R. Coulson, Proceedings of National Academy of Science U.S.A. (1977), 74, pp. 5463-5467. For example, preferably, a commercially available kit such as ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit manufactured by Perkin-Elmer can be used.

List 2

Primer 21:	ctaccaaatt ccacaactta aagttca	27mer
Primer 22:	ggaataataa gcttcacaca tactgtactc tc	32mer
15 Primer 23:	atggctccaa gctttagcaa ggaaaattcc	30mer
Primer 24:	tcaaaataag tactcaacag tggtaaaacc	30mer

The genes of the present invention derived from Cruciferae plants such as mustard (*Brassica juncea*) and rapeseed (*Brassica napus*) can be obtained by the following method.

20 For example, tissue of a Cruciferae plant such as mustard or rapeseed is frozen in liquid nitrogen and ground physically with a mortar or other means into finely divided tissue debris powder. From the tissue debris powder, RNA is extracted by a conventional method. A commercially available

RNA extraction kit can be utilized in the extraction. The RNA is recovered from thus-obtained RNA extract by ethanol precipitation. Poly-A tailed RNA is fractionated from the RNA thus recovered by a conventional method. A commercially

5 available oligo-dT column can be utilized in the fractionation.

cdNA is synthesized from the poly-A tailed RNA thus obtained by a conventional method. The synthesis can be carried out by using a commercially available cdNA synthesis kit. DNA are

10 amplified by PCR using the above-obtained cdNA as a template and primers designed and chemically synthesized on the basis of the nucleotide sequence of SEQ ID NO: 6. For example, when

PCR is carried out by using cdNA derived from mustard (*Brassica juncea*) as the template and primers 33 and 34 shown in List 3 hereinafter, the genes from Cruciferae plants of the present

15 invention, e.g., the "raffinose synthase gene having a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 5," and the "raffinose synthase gene having the nucleotide sequence represented by the 1st to 2654th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6" can be obtained.

20 According to a particular purpose, the PCR primers can also be designed and synthesized on the basis of the nucleotide sequence of SEQ ID NO: 6. For example, in order to amplify DNA encoding the open reading frame region of the "raffinose synthase gene having a nucleotide sequence encoding a protein having the

amino acid sequence of SEQ ID NO: 5", and the "raffinose synthase gene having the nucleotide sequence represented by the 134th to the 2467th nucleotides of SEQ ID NO: 6", preferably, oligonucleotides having the nucleotide sequences represented by primers 35 and 36 in List 3 are synthesized and used as the primers.

The amplified DNA can be cloned according to a conventional method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; or "Current Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc. ISBN 0-471-50338-X.

Alternatively, cloning can be carried out, for example, by using a commercially available TA cloning kit (Invitrogen) or a commercially available plasmid vector such as pBluescript II (Stratagene). The nucleotide sequence of the DNA clone can be determined by dideoxy terminating method such as that described by F. Sanger, S. Nicklen, A.R. Coulson, Proceedings of National Academy of Science U.S.A. (1977), 74, pp. 5463-5467. For example, preferably, the commercially available ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit of Perkin-Elmer can be used.

#### List 3

Primer 31:        ttggaagaga agacgccgcc gggaatcgtc    30mer  
Primer 32:        ttaagccccg gcgagagctc tggccggaca    30mer  
Primer 33:        accaatccaa aatctcatca aataatcgca    30mer

Primer 34: aaataatagg ggcagtacaa attacaccac 30mer

Primer 35: atggctccac cgagcgtaat taaatccga 29mer

Primer 36: ctaaaactca tacttaatag aagacaaacc 30mer

Then, a nucleic acid having a partial nucleotide  
5 sequence of the gene of the present invention (hereinafter  
referred to as "the gene fragment") which is obtained by the  
above-described method is labeled and then used as a probe in  
a hybridization method. The probe can be hybridized to, for  
example, DNA derived from soybean, a Chenopdiaceae plant or a  
10 Cruciferae plant to detect a nucleic acid having the probe  
specifically bound thereto, thereby detecting a nucleic acid  
having the raffinose synthase gene.

As the DNA derived from soybean, a Chenopdiaceae plant  
such as beet or a Cruciferae plant such as mustard or rapeseed,  
15 for example, a cDNA library or a genomic DNA library of these  
plants can be used. The gene library may also be a commercially  
available gene library as such or a library constructed according  
to a conventional library construction method, for example,  
described in "Molecular Cloning: A Laboratory Manual 2nd  
20 edition" (1989), Cold Spring Harbor Laboratory Press; "Current  
Protocols In Molecular Biology" (1987), John Wiley & Sons, Inc.  
ISBN 0-471-50338-X.

As the hybridization method, for example, plaque  
hybridization or colony hybridization can be employed, and they  
25 are selected depending upon the kind of vector used in the

construction of a library. More specifically, when the library to be used is constructed with a phage vector, a suitable host microorganism is mixed with the phage of the library under infectious conditions to obtain transformants. The

5 transformant is further mixed with a soft agar medium, and the mixture is plated on an agar medium. Thereafter, the mixture is cultured at 37°C until a plaque of an appropriate size appears.

When the library to be used is constructed with a plasmid vector, the plasmid is introduced into a suitable host microorganism to form transformants. The transformant obtained is diluted to a suitable concentration and the dilution is plated on an agar medium, after which it is cultured at 37°C until a colony of an appropriate size appears. In either case of the above

10 libraries, a membrane filter is placed on the surface of the agar medium after the above cultivation, so that the phage or transformant is transferred to the membrane. This membrane is denatured with an alkali, followed by neutralization, and for example, when a nylon membrane is used, the membrane is

15 irradiated with ultraviolet light, so that DNA of the phage or transformant is fixed on the membrane. This membrane is then subjected to a hybridization method wherein the gene fragment which has a partial nucleotide sequence of the gene of the present invention and labeled by a conventional method (hereinafter referred to as "the labeled gene fragment") is used as a probe.

25 For this method, reference may be made, for example, to D.M.

Glover ed., "DNA cloning, a practical approach" IRL PRESS (1985), ISBN 0-947946-18-7. There are various reagents and temperature conditions to be used in the hybridization. For example, in general, prehybridization is carried out by immersion of the  
5 membrane in a prehybridization solution [6 x SSC (0.9 M NaCl, 0.09 M citric acid), 0.1 to 1 (w/v)% SDS, 100 µg/ml denatured salmon sperm DNA] and incubation at 65°C for 1 hour. Then, hybridization is carried out by addition and mixing of the labeled gene fragment thereto and incubating the membrane at  
10 42 to 68°C for 4 to 16 hours.

In the present invention, the "stringent conditions" are those wherein incubation is carried out, for example, at 65 to 68°C in the above hybridization.

After hybridization, the membrane is taken out and  
15 is washed with 2 x SSC containing 0.1 to 1 (w/v)% SDS, further rinsed with 0.2 x SSC containing 0.1 to 1 (w/v)% SDS, and then dried. The membrane is analyzed, for example, by autoradiography or other techniques to detect the position of the probe on the membrane, thereby detecting the position on  
20 the membrane of a nucleic acid having a nucleotide sequence homologous to that of the probe used. The clone corresponding to the position of the nucleic acid thus detected on the membrane is identified on the original agar medium and the positive clone is selected so that the clone having the nucleic acid can be

isolated. The same procedures of detection are repeated to purify the clone having the nucleic acid.

Alternatively, a commercially available kit such as GENE TRAPPER cDNA Positive Selection System kit (GibcoBRL) can be used. In this method, first, a single-stranded DNA library is hybridized with the biotynylated gene fragment (i.e., probe), followed by adding streptoavidin-bound magnet beads and mixing.

From the mixture, the streptoavidin-bound magnetic beads are collected with a magnet, so that single-stranded DNA having a nucleotide sequence homologous to that of the probe used, which has been bound to these beads through the gene fragment, biotin and streptoavidin, is collected and detected. The single-stranded DNA collected can be converted into a double-strand form by reaction with a suitable DNA polymerase using a suitable oligonucleotide as a primer.

As described above, a nucleic acid containing raffinose synthase gene can obtained by detecting a nucleic acid hybridizable to the gene fragment in DNAs of a gene library derived from soybean, a Chenopdiaceae plant or a Cruciferae plant, purifying a clone having the nucleic acid and isolating phage or plasmid DNA from the clone. By preparing the restriction map or determining the nucleotide sequence of the nucleic acid thus obtained according to a conventional method, the nucleic acid containing the gene of the present invention can be confirmed.

For example, the gene of the present invention from

a Chenopodiaceae plant can be confirmed by the following point:

The amino acid encoded by the nucleotide sequence thus determined has 75% or more homology to the amino acid sequence represented by the 103rd to 208th amino acids in the amino acid  
5 sequence of SEQ ID NO: 3;

80% or more homology to the amino acid sequence represented by the 255th to 271st amino acids in the amino acid sequence of SEQ ID NO:3;

70% or more homology to the amino acid sequence  
10 represented by the 289th to 326th amino acids in the amino acid sequence of SEQ ID NO: 3; or

70% or more homology to the amino acid sequence represented by the 610th to 696th amino acids in the amino acid sequence of SEQ ID NO: 3.

15 The gene of the present invention from a Cruciferae plant can be confirmed, for example, by the following point:

The amino acid sequence encoded by the nucleotide sequence determined has 75% or more homology to the amino acid sequence represented by the 111th to 213th amino acids in the  
20 amino acid sequence of SEQ ID NO: 5;

80% or more homology to the amino acid sequence represented by the 260th to 275th amino acids in the amino acid sequence of SEQ ID NO: 5;

70% or more homology to the amino acid sequence  
25 represented by the 293rd to 325th amino acids in the amino acid



sequence of SEQ ID NO: 5; or

70% or more homology to the amino acid sequence represented by the 609th to 695th amino acids in the amino acid sequence of SEQ ID NO: 5.

5           The "homology" used herein means the proportion of the number of amino acids in a region, which are identical to those in a different region to be compared, to the number of the entire amino acids in the former region, upon comparing regions having similarity in two amino acid sequences. In this  
10       respect, it is preferred that the region having similarity contains more amino acids. Such homology of amino acid sequences can be evaluated by using a commercially available gene analysis software such as GENETIX (Software Kaihatu K.K.).

15           Further, according to the same manner as described above, a nucleic acid containing raffinose synthase gene can be detected by hybridization to DNA from the desired organism using the gene fragment as a probe to detect a nucleic acid to which the probe specifically binds (hereinafter referred to as the detection method of the present invention). The gene  
20       fragment used herein can be chemically synthesized according to a conventional method on the basis of the nucleotide sequence represented by SEQ ID NO: 2, 4, 6 or 8. Alternatively, it can be prepared by PCR using as primers oligonucleotides chemically synthesized according to a conventional method on the basis of  
25       the nucleotide sequence represented by SEQ ID NO: 2, 4, 6 or

8.

The gene fragment may be a part of the non-translated region of the raffinose synthase gene as well as the open reading frame thereof. For example, an oligonucleotide having the same nucleotide sequence as a part of that of 5'-upstream side such as the 1st to 235th nucleotides in the nucleotide sequence of SEQ ID NO: 4, the 1st to 133rd nucleotides in the nucleotide sequence of SEQ ID NO: 6 and the like, or a part of that of 3'-downstream side such as the 2588th to 2675th nucleotides in the nucleotide sequence of SEQ ID NO: 4, the 2468th to 2676th nucleotides in the nucleotide sequence of SEQ ID NO: 6 and the like.

When PCR is carried out by using the gene fragment as primers, it is possible to amplify a nucleic acid containing raffinose synthase gene from DNA derived from the desired organism (hereinafter referred to as the amplification method of the present invention).

More specifically, for example, oligonucleotides having the nucleotide sequences of the gene fragment are designed and chemically synthesized according to a conventional method.

In general, it is preferred that the number of nucleotides is more from a viewpoint that the specificity of annealing is ensured. It is, however, also preferred that the number of nucleotides is not so many from viewpoints that the primers themselves are liable to have a higher structure giving possible

deterioration of the annealing efficiency and that complicated procedures are required in the purification after the synthesis.

Normally, oligonucleotides composed of 15 to 50 bases are preferred. In this respect, based on the codon table showing the correspondence of amino acids encoded by codons, a mixture of primers can also be synthesized by using a mixture of plural bases so that a residue at a specified position in a primer is changed to different bases according to the variation of codons which can encode one certain amino acid. Alternatively, for example, a base such as inosine which can form a base pair with plural bases can be used instead of the above mixture of plural bases.

#### Coding Table

	Phe: UUU, UUC	Ser: UCU, UCC, UCA, UCG, AGU, AGC
15	Tyr: UAU, UAC	Cys: UGU, UGC
	Stop: UAA, UAG, UGA	Trp: UGG
	Leu: UUA, UUG, CUU, CUC, CUA, CUG	Pro: CCU, CCC, CCA, CCG
	His: CAU, CAC	Gln: CAA, CAG
	Arg: CGU, CGC, CGA, CGG, AGA, AGG	
20	Ile: AUU, AUC, AUA	Thr: ACU, ACC, ACA, ACG
	Asn: AAU, AAC	Lys: AAA, AAG
	Met: AUG	
	Val: GUU, GUC, GUA, GUG	Ala: GCU, GCC, GCA, GCG
	Asp: GAU, GAC	Gly: GGU, GGC, GGA, GGG
25	Glu: GAA, GAG	

In the above codon table, each codon is shown as the nucleotide sequence in mRNA and its right hand is the 5'-terminus.

U represents uracil base in RNA and corresponds to thymine base in DNA.

5           An oligonucleotide having the same nucleotide sequence as the coding strand of the double-stranded DNA of the gene of the present invention is called a "sense primer" and that having a nucleotide sequence complementary to the coding strand is called an "antisense primer".

10           A sense primer having the same nucleotide sequence as that of 5'-upstream side in the coding strand of the gene of the present invention, and an antisense primer having a nucleotide sequence complementary to the nucleotide sequence on the 3'-downstream side in this coding strand are used in  
15 combination for PCR reaction, for example, with a gene library, genomic DNA or cDNA as the template to amplify DNA. As the gene library to be used, for example, there are a cDNA library and a genomic library derived from soybean, a Chenopodiaceae plant such as beet or a Cruciferae plant such as mustard or rapeseed,  
20 etc. The gene library may also be a library constructed according to a conventional library construction method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor Laboratory Press; "Current Protocols In Molecular Biology" (1987), John Wiley &  
25 Sons, Inc. ISBN 0-471-50338-X, or a commercially available gene

library as such. As the genomic DNA or cDNA, for example, there are those prepared from soybean, a Chenopdiaceae plant such as beet or a Cruciferae plant such as mustard or rapeseed, etc.

For example, PCR is carried out by using the primers 31 and 32 in the above List 3 and as the template cDNA derived from mustard to amplify DNA having the nucleotide sequence

represented by the 749th to 1215th nucleotides in the nucleotide sequence of SEQ ID NO: 6. Further, PCR is carried out by using the primers and as the template cDNA derived from rapeseed to amplify DNA having the nucleotide sequence represented by the 1st to 467th nucleotides in the nucleotide sequence of SEQ ID NO: 8. The nucleic acid thus amplified can be confirmed by conventional electrophoresis. The nucleic acid can be cloned according a conventional method such as that described in

"Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor laboratory Press or "Current Protocols in Molecular Biology" (1987), John Wiley & Sons, Inc., ISBN 0-471-50338-X. For the nucleic acid, its restriction map is prepared or its nucleotide sequence is determined by a

conventional method, so that the nucleic acid containing raffinose synthase gene or a part thereof can be identified.

When the nucleic acid contains a part of raffinose synthase, PCR can be carried out on the basis of its nucleotide sequence to amplify the nucleic acid containing the 5'-upstream side nucleotide sequence or the 3'-downstream side nucleotide

sequence. That is, based on the nucleotide sequence of the above-obtained nucleic acid, an antisense primer is designed and synthesized for amplification of the 5'-upstream side part, and a sense primer is designed and synthesized for amplification of the 3'-downstream side part. The nucleotide sequence of the 5'-upstream side part or 3'-downstream side part of the nucleotide sequence already obtained can be determined by the RACE method using these primers and a commercially available kit such as Marathon Kit of Clontech. The full length raffinose synthase gene can be obtained by synthesizing new primers based on both terminal sequences in the nucleotide sequence thus determined and carrying out PCR again.

The above detection method of the present invention can also be used in the analysis of genotypes of a plant such as soybean, a Chenopodiaceae plant or a Cruciferae plant, etc.

More specifically, for example, a genomic DNA derived from soybean, a Chenopodiaceae plant or a Cruciferae plant is prepared according to a conventional method, for example, described in "Cloning and Sequence (Plant Biotechnology Experiment Manual)" compiled under the supervision of Itaru Watanabe, edited by Masahiro Sugiura, published by Noson Bunka-sha, Tokyo (1989).

The genomic DNA is digested with at least several kinds of restriction enzymes, followed by electrophoresis. The electrophoresed DNA is blotted on a filter according to a conventional method. This filter is subjected to hybridization

with a probe prepared from DNA having the gene fragment by a conventional method, and DNA to which the probe hybridizes is detected. The DNAs detected are compared in length between different varieties of a specified plant species. The

5 differences in length make possible the analysis of differences in phenotypic characteristics accompanied with the expression of raffinose family oligosaccharides between these varieties.

Furthermore, when the DNAs detected by the above method are compared in length between the gene recombinant plant and the  
10 non-gene recombinant plant of the same variety, the former plant can be distinguished from the latter plant by the detection of hybridizing bands greater in number or higher in concentration for the former plant than for the latter plant. This method can be carried out according to the RFLP (restriction fragment length  
15 polymorphism) method, for example, described in "Plant PCR Experiment Protocols" compiled under the supervision of Ko Shimamoto and Takuji Sasaki, published by Shujun-sha, Tokyo (1995), ISBN 4-87962-144-7, pp. 90-94.

Further, the amplification method of the present  
20 invention can be used for an analysis of genes of soybean, a Chenopdiaceae plant or a Cruciferae plant, etc. More specifically, for example, the amplification method of the present invention is carried out by using plant genomic DNA prepared from soybean, a Chenopdiaceae plant or a Cruciferae  
25 plant to amplify DNA. The amplified DNA is mixed with a

formaldehyde solution, followed by heat denaturing at 85°C for 5 minutes and then quickly cooling on ice. A sample thereof is subjected to electrophoresis on, for example, 6 (w/v)% polyacrylamide gel containing 0 (v/v)% or 10 (v/v)% of glycerol.

5 For this electrophoresis, a commercially available electrophoresis apparatus such as that for SSCP (Single Strand Conformation Polymorphism) can be used and the electrophoresis can be carried out with maintaining the gel at a constant temperature, for example, at 5°C, 25°C, 37°C, etc. From the  
10 electrophoresed gel, DNA is detected, for example, by a method such as silver staining method with a commercially available reagent. From the differences of behavior between the varieties in the electrophoresis of the DNA detected, a mutation in the raffinose synthase gene is detected, and an analysis is carried  
15 out for differences caused by the mutation in phenotypic characteristics accompanied with the expression of raffinose family oligosaccharides. This method can be carried out according to the SSCP method, for example, described in "Plant PCR Experiment Protocols" compiled under the supervision of Ko  
20 Shimamoto and Takuji Sasaki, published by Shujun-sha, Tokyo (1995), ISBN 4-87962-144-7, pp. 141-146.

The analysis of the plant gene from soybean, a Chenopodiaceae plant or a Cruciferae plant by the above detection method or amplification method of the present invention can be  
25 used not only for the analysis of differences in phenotypic



characteristics accompanied with the expression of raffinose family oligosaccharides, but also, for example, for the selection of clones having the desired characters upon production of a novel variety of soybean, a Chenopdiaceae plant or a Cruciferae plant. Further, it can also be used for identification of a clone thus produced and having the characters derived from a recombinant plant upon producing a plant variety using the recombinant plant.

For expression of the gene of the present invention in cells of a host, preferably, a nucleic acid comprising a nucleic acid fragment which contains the gene of the present invention, and a nucleic acid fragment which has a promoter activity in the host cells and joined to the former nucleic acid fragment (hereinafter referred to as the expression nucleic acid of the present invention) can be used.

The nucleic acid fragment having promoter activity in the expression nucleic acid of the present invention is not limited to a specific one, so long as it is functionable in a host to be transformed. For example, there are synthetic promoters functionable in *Escherichia coli*, such as *E. coli* lactose operon promoter, *E. coli* tryptophan operon promoter and tac promoter, etc.; yeast alcohol dehydrogenase gene (ADH) promoter, adenovirus major late (Ad.ML) promoter, SV40 early promoter, baculovirus promoter and the like. When the host is

a plant, the promoter may include, for example, T-DNA derived constitutive promoters such as nopaline synthase gene (NOS) promoter, octopine synthase gene (OCS) promoter, etc.; plant virus-derived promoters such as cauliflower mosaic virus

5 (CaMV)-derived 19S and 35S promoters; inducible promoters such as phenylalanine ammonia-lyase (PAL) gene promoter, chalcone synthase (CHS) gene promoter, pathogenesis-related protein (PR) gene promoter, etc. Furthermore, vector pSUM-GY1 (see JP-A 06-189777/1994) can also be used, which has a promoter giving  
10 specific expression in a specified plant tissue, e.g., a promoter of soybean-derived seed storage protein glycinin gene (JP-A 6-189777).

Furthermore, a nucleic acid fragment having a terminator activity can be joined to the expression nucleic acid  
15 of the present invention. In this case, it is generally preferred that the expression nucleic acid of the present invention is constructed so that the nucleic acid fragment having a terminator activity is positioned downstream the raffinose synthase gene. The terminator to be used is not particularly  
20 limited, so long as it is functionable in cells of a host to be transformed. For example, when the host is a plant, there are T-DNA derived constitutive terminators such as nopaline synthase gene (NOS) terminator, etc.; plant derived terminators such as terminators of allium virus GV1 or GV2, and the like.

25 The expression nucleic acid of the present invention

can be introduced into a host cell according to a conventional gene engineering technique to obtain a transformant. If necessary, the expression nucleic acid of the present invention can be inserted into a vector having a suitable marker depending upon a particular transformation technique for introduction of the nucleic acid into a host cell.

A vector into which the expression nucleic acid of the present invention is inserted can be introduced into a microorganism according to a conventional method, for example, described in "Molecular Cloning: A Laboratory Manual 2nd edition" (1989), Cold Spring Harbor laboratory Press or "Current Protocols in Molecular Biology" (1987), John Wiley & Sons, Inc., ISBN 0-471-50338-X. The microorganism transformed with the vector can be selected on the basis of a selection marker such as antibiotic resistance, auxotrophy or the like. In case that the gene of the present invention is joined to the downstream of an inducible promoter (e.g., tac promoter) in the translatable form in the selected microorganism (e.g., *E. coli* transformant), a translated product of the gene of the present invention can be expressed under conventional culture and inducible conditions and can be recovered as a peptide or a protein.

The raffinose synthase activity of the translated product of the gene of the present invention thus prepared can be measured by, for example, a method described in L. Lehle and W. Tanner, Eur. J. Biochem., 38, 103-110 (1973) to identify the

translated product having the "capability of binding D-galactosyl group through  $\alpha$  (1 $\rightarrow$ 6) bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in sucrose molecule". More specifically, for example,

5 the gene of the present invention is cloned in pGEX-4T3 (Pharmacia) to obtain a plasmid containing the expression nucleic acid of the present invention. The resultant plasmid is introduced into, for example, *E. coli* HB101 strain to obtain a transformant. The resultant transformant is culture

10 overnight and 1 ml of the culture is inoculated into 100 ml of LB culture medium. It is incubated at 37°C for about 3 hours and IPTG (isopropylthio- $\beta$ -D-galactoside) is added at a final concentration of 1 mM, followed by further incubation for 5 hours.

Cells are recovered from the culture broth by centrifugation

15 and are suspended by addition of 10 times of the cell weight of 100 mM Tris-HCl (pH 7.4), 1 mM EDTA, 5 mM DTT (dithiothreitol), 1 mM PMSF (phenylmethylsulfonyl fluoride) and 1 mM benzamide.

The suspension is sonicated with a ultrasonic disrupter (Branson) to disrupt the cells. The disrupted cell suspension

20 is centrifuged to recover a soluble protein solution. The resultant protein solution is added to a reaction mixture containing at final concentrations of 100 mM Tris-HCl (pH 7.4), 5 mM DTT (dithiothreitol), 0.01% BSA, 200  $\mu$ M sucrose, 5 mM galactinol and 31.7  $\mu$ M [ $^{14}$ C] sucrose. The reaction mixture is

25 incubated at 37°C, followed by addition of 1.5 times in volume

of ethanol and stirring. Insoluble materials are removed by centrifugation, the supernatant is spotted on, for example, a HPTLC cellulose thin layer chromatography plate (Merck, HPTLC plates cellulose) and then the plate is developed with n-butanol-pyridine-water-acetic acid (60 : 40 : 30 : 3). The developed plate is dried and analyzed with an imaging analyzer (FUJIX Bio-Image Analyzer BAS-2000II manufactured by Fuji Film) to determine [<sup>14</sup>C] raffinose produced to measure the raffinose synthase activity.

In addition, the translated product as prepared above can also be used as an antigen for producing an antibody. The antibody thus produced can be used for, for example, detection and determination of the gene of the present invention in a crude protein extract prepared from an organism such as a plant.

When the host is a plant, the vector into which the gene of the present invention is inserted can be introduced into plant cells by a conventional means such as *Agrobacterium* infection method (JP-B 2-58917 and JP-A 60-70080), electroporation into protoplasts (JP-A 60-251887 and JP-B 5-68575) or particle gun method (JP-A 5-508316 and JP-A 63-258525). The plant cell transformed by the introduction of the vector can be selected on the basis of a selection marker, for example, resistance to an antibiotic such as kanamycin or hygromycin. From the plant cell thus transformed, a transformed plant can be regenerated by a conventional plant cell cultivation

method, for example, described in "Plant Gene Manipulation Manual (How to Produce Transgenic Plants)" written by Uchimiya, 1990, Kodan-sha Scientific (ISBN 4-06-153513-7), pp. 27-55.

Furthermore, the collection of seeds from the transformed plant also makes it possible to proliferate the transformed plant. In addition, crossing between the transformed plant obtained and the non-transformed plant makes it possible to produce progenic plants with the characters of the transformed plant.

As gene engineering techniques in soybean, basically, the above general techniques can be employed. More specifically, "transformation of soybean plant strain by particle gun" described in EP 301749, gene introduction methods. for example, described in Torisky, R.S., Kovacs, L., Avdiushko, S., Newman, J.D., Hunt, A.G. and Collins, G.B., "Development of a binary vector system for plant transformation based on the supervirulent *Agrobacterium tumefaciens* strain Chry5", Plant Cell Rep., (1997), 17, p. 102-108, etc. can be employed.

As gene engineering techniques in a *Chenopodiaceae* plant, basically, the above general techniques can be employed.

More specifically, gene introduction methods, for example, described in M. Mannerlof, S. Tuveesson, P. Steen and P. Tenning, "Transgenic sugar beet tolerant to glyphosate", *Euphytica* (1997), 94, p 83-91, B.K. Konwar, "Agrobacterium tumefaciens-Mediated Genetic Transformation of Sugar Beet (*Beta vulgaris* L.)", *J. Plant Biochemistry & Biotechnology*

(1994), 3, p. 37-41 can be employed.

As gene engineering techniques in a Cruciferae plant, basically, the above general techniques can be employed. More specifically, the gene introduction can be carried out according to a method, for example, described in J. Fry, A. Barnason and R.B. Horsch, "Transformation of Brassica napus with *Agrobacterium tumefaciens* based vectors", Plant Cell Reports (1987), 6, 321-325.

For example, when gene introduction is carried out by *Agrobacterium* infection method, first, the above-described expression nucleic acid of the present invention is inserted into a binary vector. The resultant vector can be introduced into, for example, *Agrobacterium tumefaciens* LBA 4404 strain which has been converted into a competent state by treatment with calcium chloride. A transformant can be selected by an appropriate selection method according to the selection marker gene of the vector, for example, cultivation of a strain containing the vector in a culture medium containing an antibiotic in case that the selection marker gene is that giving resistance to the antibiotic such as kanamycin. The resultant transformed *Agrobacterium* strain can be culture in a liquid culture medium, for example, LB medium.

Soybean, a Chenopodiaceae plant or a Cruciferae plant can be transformed by using thus obtained *Agrobacterium*

transformant culture broth as described below. For example, seeds from soybean, beet, rapeseed or mustard is sowed aseptically in, for example, 1/2 MS medium containing 2% sucrose and 0.7% agar. After about 1 week, cotyledons and petioles of the germinated plant are cut off with a scalpel aseptically and transplanted in, for example, MS medium containing 3% sucrose, 0.7% agar, 4.5  $\mu\text{M}$  BA, 0.05  $\mu\text{M}$  2,4-D and 3.3  $\mu\text{M}$   $\text{AgNO}_3$  and cultured for one day. The cotyledons and petioles thus precultured are transferred to 1000-fold dilution of the above *Agrobacterium* culture broth and allowed to stand for 5 minutes. The cotyledons and petioles are transferred to the same medium as that of the preculture again and cultured for about 3 to 4 days. The cotyledones and petioles thus cultured are transferred to, for example, MS medium containing 3% sucrose, 4.5  $\mu\text{M}$  BA, 0.05  $\mu\text{M}$  2,4-D, 3.3  $\mu\text{M}$   $\text{AgNO}_3$  and 500 mg/liter cefotaxim, followed by shaking for 1 day to remove microbial cells. The resultant cotyledons and petioles are transferred to, for example, MS medium containing 3% sucrose, 0.7% agar, 4.5  $\mu\text{M}$  BA, 0.05  $\mu\text{M}$  2,4-D, 3.3  $\mu\text{M}$   $\text{AgNO}_3$  100 mg/liter cefotaxim and 20 mg/liter kanamycin, followed by culturing for 3 to 4 weeks. Then, the cotyledons and petioles are transferred to, for example, MS medium containing 3% sucrose, 0.7% agar, 4.5  $\mu\text{M}$  BA, 0.05  $\mu\text{M}$  2,4-D, 100 mg/liter cefotaxim and 20 mg/liter kanamycin and cultured. Culture in this medium is continued with subculturing every 3 to 4 weeks. When a shoot are regenerated, it is subcultured in,



for example, MS medium containing 3% sucrose, 0.7% agar and 20 mg/liter kanamycin for 3 to 4 weeks. When the plant makes roots, it is transferred to vermiculite-peat moss (1 : 1) and acclimatized by culturing at 21 to 22°C under day and night conditions of 12 hours: 12 hours = day time : night. As the plant grows, it is transferred to appropriate cultivation soil to culture the plant. A genomic DNA is extracted from the leaf of the regenerated plant according to the above method and PCR is carried out by using as primers having partial nucleotide sequences of the expression nucleic acid of the present invention to confirm the insertion of the gene of the present invention into the plant.

As described hereinabove, by introducing the gene of the present invention into a plant, for example, soybean, a Chenopdiaceae plant or a Cruciferae plant, it is possible to vary the expression level and activity of raffinose synthase in the plant to control the amount of raffinose family oligosaccharides in the plant. The gene of the present invention is useful in techniques for varying the expression level and activity of raffinose synthase in soybean, a Chenopdiaceae plant or a Cruciferae plant on the basis of gene homology, for example, techniques such as homologous recombination and antisense technique, cosuppression and the like.

The following examples further illustrate the present

invention in detail but are not to be construed to limit the scope of the present invention.

Example 1

Preparation of cDNA Derived from Soybean

5           About 2 g of immature seeds of soybean (*Glycine max*) Williams82 were frozen in liquid nitrogen and then ground with a mortar, to which 20 ml of Isogen (Nippon Gene) was added, and the mixture was further thoroughly ground. The ground material was transferred into a centrifugation tube, to which 4 ml of  
10 chloroform was added, and the mixture was stirred with a vortex mixer and then centrifuged at 6,500 x g for 10 minutes at 4°C.

The water layer was collected, to which 10 ml of isopropanol was added, and the mixture was stirred and then centrifuged at 6,500 x g for 10 minutes at 4°C. The resulting precipitate was washed  
15 with 10 ml of 70% ethanol and then dissolved in 1 ml of elution buffer (10 mM Tris-HCl/pH 7.5, 1 mM EDTA, 0.1% SDS). The solution was allowed to stand at 60°C for 10 minutes and then centrifuged at 10,000 x g for 1 minute to remove insoluble matter. To the resulting supernatant was added an equivalent volume of  
20 Oligotex-dT30 (Takara), and the mixture was stirred and then allowed to stand at 65°C for 5 minutes. Further, the mixture was placed on ice and allowed to stand for 3 minutes, to which 200 µl of 5 M NaCl was added, and the mixture was mixed and then allowed to stand at 37°C for 10 minutes. The mixture was then centrifuged

at 10,000 x g for 3 minutes at 4°C. The precipitate was collected and then suspended in 1 ml of TE buffer, and the suspension was allowed to stand at 65°C for 5 minutes. Further, the suspension was placed on ice and then allowing to stand for 3 minutes, followed  
5 by centrifugation at 10,000 x g for 3 minutes at 4°C to remove precipitate.

To the resulting supernatant were added 100 µl of 3M sodium acetate and 2 ml of ethanol to precipitate and collect RNA. The collected RNA was washed twice with 70% ethanol and then  
10 dissolved in 20 µl of sterilized water, which was used for the subsequent cDNA synthesis. The amount of RNA obtained was determined by the measurement of absorbance at 260 nm.

For the cDNA synthesis, First Strand Synthesis Kit for RT-PCR (Amersham) and cDNA Synthesis Kit (Takara) were used, and  
15 all operations were made according to the protocol attached to kits.

#### Example 2

##### Cloning of Raffinose Synthase Gene from Soybean cDNA

PCR was carried out by using the cDNA obtained from  
20 immature seeds of soybean (*Glycine max*) Williams82 in Example 1 as a template and the primers designed on the basis of the amino acid sequence of SEQ ID No: 1, i.e., primers having nucleotide sequences shown in List 4 below to amplify a DNA fragment. The PCR was carried out with Gene Amp PCR Systems 2400 and DNA Thermal

Cycler Model 480 of Perkin-Elmer using Advantage KlenTaq cDNA Kit of Clontech. The reaction was carried out by repeating the cycle for maintaining at 94°C for 1 minute, at 50°C for 3 minutes and then at 72°C for 3 minutes 40 times to amplify the DNA fragment.

5 The amplified DNA fragment was cloned with TA cloning kit (Invitrogen), followed by sequence reaction using ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit of Perkin-Elmer and analysis of the nucleotide sequence with a 373S DNA sequencer of ABI. Based on this sequence, primers having nucleotide  
10 sequences shown in List 5 below were synthesized. The synthesis of cDNA was carried out with Marathon Kit of Clontech using mRNA obtained from leaves of soybean Williams82 in Example 1. The cDNA obtained was ligated to an adapter contained in the kit with ligase. These operations were carried out according to the protocol  
15 attached to the kit. By using the adapter-ligated cDNA thus prepared, PCR was carried out with the primers shown in List 5 according to the same manner as the above. The nucleotide sequence in terminal region of the gene was analyzed according to the protocol attached to the Marathon Kit of Clontech. As a result,  
20 the nucleotide sequence of SEQ ID NO: 2 was determined.

List 4

4-5-F primer:

cgatggatgg giaaittiat icaiccigai tgggaiatgt t 41mer

4-6-RV primer:

ggccacatit tiacia(ag)icc iatiggigci aa 32mer

List 5

5-SC-2:

tgttactagg cgaaacaaga gtagctctga 30mer

5 Example 3

Preparation of cDNA derived from Chenopdiaceae Plant

About 2 g of leaves of beet (*Beta vulgaris*: haming) was frozen in liquid nitrogen and then ground with a mortar, to which 20 ml of Isogen (Nippon Gene) was added, and the mixture was further thoroughly ground. The ground material was transferred into a centrifugation tube, to which 4 ml of chloroform was added, and the mixture was stirred with a vortex mixer and then centrifuged at 6,500 x g for 10 minutes at 4°C. The water layer was collected, to which 10 ml of isopropanol was added, and the mixture was stirred and then centrifuged at 6,500 x g for 10 minutes at 4°C. The resulting precipitate was washed with 10 ml of 70% ethanol and then dissolved in 180 µl of DEPC-treated sterilized water. The solution was allowed to stand at 55°C for 5 minutes and 20 µl of 5 M NaCl was added thereto. The resulting solution was purified using BIOMAG mRNA PURIFICATION KIT (PerSeptive Biosystems: Catalog No. 8-MB4003K).

To the resulting mRNA solution were added 3M sodium acetate and ethanol, and RNA was precipitated and collected. The collected RNA was washed twice with 70% ethanol and then dissolved

in 20 µl of sterilized water, which was used for the subsequent cDNA synthesis. The amount of RNA obtained was determined by the measurement of absorbance at 260 nm.

For the cDNA synthesis, SMART PCR cDNA Synthesis Kit (Clontech) was used, and all operations were made according to the protocol attached to the kit.

#### Example 4

Analysis of Nucleotide Sequence of Raffinose Synthase Gene from Chenopdiaceae Plant

Synthetic DNA primers having the nucleotide sequences shown in List 6 below were synthesized. The PCR method was carried out with Gene Amp PCR Systems 2400 and DNA Thermal Cyclor Model 480 of Perkin-Elmer using Advantage KlenTaq cDNA Kit of Clontech.

The PCR was carried out with the above primers and cDNA of beet obtained in the above Example 3 by repeating the cycle for maintaining at 94°C for 1 minute, at 50°C for 3 minutes and then at 72°C for 3 minutes 40 times. As a result, the combinations of primers 6-3-F and 6-8-RV and primers 6-10-F and 6-6-RV gave an amplification of about 0.3 kb and 0.6 kb bands, respectively. The amplified DNA fragments were cloned with TA cloning kit

(Invitrogen), followed by sequence reaction using ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit of Perkin-Elmer and analysis of nucleotide sequence with a 373S DNA sequencer of ABI. Based on the resulting nucleotide sequences, synthetic DNA

primers having nucleotide sequences shown in List 7 below were prepared and PCR was carried out using cDNA obtained from beet in Example 3 in the same manner as above. As a result, DNA having the nucleotide sequence of SEQ ID NO: 4 was finally obtained from cDNA of beet.

List 6

6-3-F:

cgaggigggt gicciccigg ittigtati atigaigaig gitggca 47mer

6-8-RV:

at(t/c)tt(a/g)tcia cigcia(a/g) (a/g)tc (t/c)tccatigt 29mer

6-10-F:

ggiacitait gg(c/t)ticaigg itgicaiatg gticaigt 38mer

6-6-RV:

ggccacatit tiacia(a/g)icc iatiggigci aa 32mer

List 7

7-Sb-1:

atctatttgt catgacgatg atccga 26mer

7-Sb-2RV:

ggccctcatt cccatattgg gatgatcctc 30mer

7-Sb-3RV:

aagcatgcc aacatacaca tgctcaacag 30mer

7-Sb-4RV:

agaccgggg aaagctttgg ggttactact 30mer

7-Sb-5:

tggatgggaa actttataca ccctgact 28mer

7-Sb-6:

gacatgttcc catctacaca cccttggtg 28mer

7-Sb-7:

ccaatttatg ttagtgatgt tgttggcaag 30mer

5 7-Sb-8RV:

tgcactccca gggtagaatt gtcatc 26mer

Example 5

Preparation of cDNA Derived from Cruciferae Plant

About 2 g of leaves of mustard (*Brassica juncea*) was  
10 frozen in liquid nitrogen and then ground with a mortar, to which  
20 ml of Isogen (Nippon Gene) was added, and the mixture was  
further thoroughly ground. The ground material was transferred  
into a centrifugation tube, to which 4 ml of chloroform was added,  
and the mixture was stirred with a vortex mixer and then  
15 centrifuged at 6,500 x g for 10 minutes at 4°C. The water layer  
was collected, to which 10 ml of isopropanol was added, and the  
mixture was stirred and then centrifuged at 6,500 x g for 10 minutes  
at 4°C. The resulting precipitate was washed with 10 ml of 70%  
ethanol and then dissolved in 180 µl of DEPC-treated sterilized  
20 water. The solution was allowed to stand at 55°C for 5 minutes  
and to which 20 µl of 5 M NaCl was added. The resulting solution  
was purified using BIOMAG mRNA PURIFICATION KIT (PerSeptive  
Biosystems: Catalog No. 8-MB4003K).

To the resulting mRNA solution were added 3M sodium



acetate and ethanol, and RNA was precipitated and collected. The collected RNA was washed twice with 70% ethanol and then dissolved in 20 µl of sterilized water, which was used for the subsequent cDNA synthesis. The amount of RNA obtained was determined by the measurement of absorbance at 260 nm.

For the cDNA synthesis, SMART PCR cDNA Synthesis Kit (Clontech) was used, and all operations were carried out according to the protocol attached to the kit.

In the same manner as described in the above, mRNA was purified from immature seeds of rapeseed Westar (*Brassica napus*) and cDNA was synthesized.

#### Example 6

Isolation and Nucleotide Sequence Analysis of Raffinose Synthase Gene derived from Cruciferae Plant

DNA primers having the nucleotide sequences shown in List 8 below were synthesized. PCR was carried out with Gene Amp PCR Systems 2400 and DNA Thermal Cycler Model 480 of Perkin-Elmer using Advantage KlenTaq cDNA Kit of Clontech. The PCR was carried out with the above primers and cDNA of mustard obtained in Example 5 by repeating the cycle for maintaining at 94°C for 1 minute, at 50°C for 3 minutes and then at 72°C for 3 minutes 40 times. The reaction products were analyzed by agarose gel electrophoresis. As a result, an amplification of about the 1.2 kb bands was detected. The amplified DNA fragment was cloned with TA cloning kit

(Invitrogen), followed by sequence reaction using ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction Kit of Perkin-Elmer and nucleotide sequence analysis with a 373S DNA sequencer of ABI.

Based on the resulting nucleotide sequence, synthetic primers  
5 having the nucleotide sequences shown in List 9 below were prepared and PCR was carried out using cDNAs from mustard (*Brassica juncea*) and rapeseed Westar (*Brassica napus*) obtained in Example 5 according to the same manner as the above. As a result, the nucleotide sequence represented by the 749th to 1215th nucleotides  
10 of SEQ ID NO: 6 and by the 1st to 467th nucleotides of SEQ ID NO: 8 were finally determined for cDNA from mustard (*Brassica juncea*) and rapeseed Westar (*Brassica napus*), respectively.

List 8

8-#1:  
15 cgattiaaig titggtggac iacicaitgg gtigg 35mer  
8-#10RV:  
caitgiacca titgicaicc itgia(ag)ccai taigticc 38mer

List 9

9-primer-1:  
20 gttagggttc atatgaacac cttcaagctc 30mer  
9-primer-2RV:  
caacggcgag atcttgcac gtcaac 26mer

Example 7

Nucleotide Sequence Analysis of Raffinose Synthase

# Full-Length Gene Derived from Cruciferae Plant

Based on the nucleotide sequences obtained in Example 6, DNA primers having the nucleotide sequences shown in List 10 below were synthesized. The cDNAs from mustard (*Brassica juncea*) and rapeseed Westar (*Brassica napus*) obtained in the same manner as described in Example 5 were ligated to adapters contained in Marathon Kit of Clontech. By using the adapter-ligated cDNAs thus prepared, PCR was carried out with primers shown in List 10. 10-B-2RV, 10-B-3RV and 10-B-4RV primers were used for nucleotide analysis of 5'-termini, and 10-B-1, 10-B-8, 10-B-7 and 10-B-6 primers were used for nucleotide analysis of 3'-termini. The nucleotide sequences were analyzed according to the protocol attached to the Marathon Kit of clontech. As a result, the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 8 were determined from mustard (*Brassica juncea*) and rapeseed Westar (*Brassica napus*), respectively.

## List 10

10-B-2RV:  
ggattcgaca caaaccgcca cgtcatcgtc 30mer  
20 10-B-3RV:  
ccacgtgcac caccggaact tatcgac 27mer  
10-B-4RV:  
aacatcgata ccatcggagt catgtccaat 30mer  
10-B-1:

gttagggttc atatgaacac cttcaagctc 30mer

10-B-8:

tctacgtctg gcacgcgctt tgcggctac 29mer

10-B-7:

5 gttgacgtca tccacatatt ggagatgttg t 31mer

10-B-6:

gttatcgcta gcatggagca ctgtaatga 29mer

#### Example 8

### 10 Construction of Expression Vectors in Plant for Raffinose Synthase Gene Derived from Cruciferae Plant

Based on the nucleotide sequence of raffinose synthase gene from mustard obtained in Example 7, DNA primers having the nucleotide sequences shown in List 11 were prepared. By using cDNA of mustard, PCR was carried out in the same manner as

15 described in Example 6. The amplified DNA fragment was digested with SacI. The DNA fragment thus digested was ligated to the vector pBI121(-) previously digested with SacI by using Ligation Kit (Takara). Plasmid pBI121 (Clontech) were digested with BamHI and SacI, and ligated to linkers shown in List 12 to prepare

20 the vector pBI121(-). The vector thus obtained was analyzed by a restriction map and PCR using primers having nucleotide sequences shown in List 13, and confirmed the direction of inserted raffinose synthase gene. The vector whose raffinose synthase gene from mustard was inserted in the expressible

25 direction was designated BjRS-Sac(+)-121 and the one whose

raffinose synthase gene from mustard was inserted in the reverse direction was designated BjRS-Sac(-)-121.

List 11

11-SacI-BjN:

5 aacgagctca atccaaaatc tcatcaaata atcgc 35mer

11-SacI-BjintRV:

acaatagttg agggcggaag agtag 25mer

List 12

12-BamSac-(+)linker:

10 gatcgagctc gtgtcggatc cagct 25mer

12-BamSac-(-)linker:

ggatccgaca cgagctc 17mer

List 13

13-35S-3:

15 cctcctcgga ttccattgcc cagctatctg 30mer

13-B-2RV:

ggattcgaca caaaccgcca cgtcatcgtc 30mer

13-B-8:

tctacgtctg gcacgcgctt tgcggctac 29mer

20 Example 9

Transformation with Raffinose Synthase Gene Derived from Cruciferae Plant

The vectors BjRS-Sac(+)-121 and BjRS-Sac(-)-121 prepared in Example 8 were used for the transformation of mustard (Brassica juncea) by the Agrobacterium infection method.

*Agrobacterium tumefaciens* (strain LBA4404 having rifampicin and streptomycin resistance) previously converted into a competent state by calcium chloride treatment was transformed independently with two plasmids BjRS-Sac(+)-121 and BjRS-Sac(-)-121 prepared in Example 8. The transformants were selected on LB medium containing 50 µg/ml rifampicin and 25 µg/ml kanamycin by utilizing the kanamycin resistant character conferred by the kanamycin resistant gene (neomycin phosphotransferase, NPTII) of the introduced plasmids.

The transformant *Agrobacterium* obtained (*Agrobacterium tumefaciens* strain LBA4404: rifampicin and streptomycin resistant) was cultured on LB medium containing 50 µg/ml rifampicin and 25 µg/ml kanamycin at 28°C for a whole day and night, and the culture was used for the transformation of mustard by the method described below.

The seeds of mustard were aseptically sowed on 1/2 MS medium containing 2% sucrose and 0.7% agar. After one week, cotyledons and petioles of sprouting plants were cut out with a scalpel, and transferred to MS medium containing 3% sucrose, 0.7% agar, 4.5 µM BA, 0.05 µM 2,4-D and 3.3 µM AgNO<sub>3</sub>, followed by preculture for 1 day. The precultured cotyledons and petioles were transferred in a 1000-fold dilution of the *Agrobacterium* culture broth and allowed to stand for 5 minutes.

The cotyledons and petioles were transferred again to the same

medium as used in the preculture, and cultured for 3 to 4 days.

The cultured cotyledons and petioles were transferred to MS medium containing 3% sucrose, 4.5  $\mu$ M BA, 0.05  $\mu$ M 2.4-D, 3.3  $\mu$ M AgNO<sub>3</sub> and 500 mg/l cefotaxim, and shaken for 1 day to remove microbial cells. The cotyledons and petioles thus treated were transferred to MS medium containing 3% sucrose, 0.7% agar, 4.5  $\mu$ M BA, 0.05  $\mu$ M 2.4-D, 3.3  $\mu$ M AgNO<sub>3</sub>, 100 mg/l cefotaxim and 20 mg/l kanamycin, and cultured for 3 to 4 weeks. The cotyledons and petioles were transferred to MS medium containing 3% sucrose, 0.7% agar, 4.5  $\mu$ M BA, 0.05  $\mu$ M 2.4-D, 100 mg/l cefotaxim and 20 mg/l kanamycin, and cultivated. The cultivation on this medium was continued with subculturing at intervals of 3 to 4 weeks.

When shoots are began to regenerate, these shoots are subcultured on MS medium containing 3% sucrose, 0.7% agar and 20 mg/l kanamycin, and cultivated for 3 to 4 weeks. The rooting plants are transferred to vermiculite : peat moss = 1 : 1, and cultivated at 21°C to 22°C in a cycle of day/night = 12 hours : 12 hours. With the progress of plant body growth, the plants are grown with cultivation soil.

#### BRIEF DESCRIPTION OF SEQUENCES

SEQ ID NO: 1 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene of the present invention.

SEQ ID NO: 2 shows a nucleotide sequence of the raffinose synthase gene of the present invention.

SEQ ID NO: 3 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene obtained from beet.

SEQ ID NO: 4 shows a cDNA nucleotide sequence of the raffinose synthase gene obtained from beet.

SEQ ID NO: 5 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene obtained from mustard.

SEQ ID NO: 6 shows a cDNA nucleotide sequence of the raffinose synthase gene obtained from mustard.

SEQ ID NO: 7 shows an amino acid sequence of a raffinose synthase protein encoded by the raffinose synthase gene obtained from rapeseed.

SEQ ID NO: 8 shows a cDNA nucleotide sequence of the raffinose synthase gene obtained from rapeseed.

List 1:

The nucleotide sequences shown in List 1 are examples of the typical primers used in the amplification of a DNA fragment having a raffinose synthase gene. All of these sequences are based on the nucleotide sequence of SEQ ID NO: 2. Primer 11 is a sense primer and Primer 12 is an antisense primer. Depending upon the purpose, recognition sequences for suitable restriction enzymes can be added to the 5'-termini of these nucleotide sequences.



List 2:

The nucleotide sequences shown in List 2 are examples of the typical primers used in the amplification of a cDNA of a raffinose synthase gene. Primer 21 is a sense primer

5 corresponding to the 5'-terminus of the beet-derived raffinose synthase gene. Primer 22 is an antisense primer corresponding to the 3'-terminus. Depending upon the purpose, recognition sequences for suitable restriction enzymes can be added to the 5'-termini of these nucleotide sequences.

10 Primer 23 is a sense primer corresponding to the N-terminus of the open reading frame, and primer 24 is an antisense primer corresponding to the C-terminus.

List 3:

15 Among the nucleotide sequences shown in List 3, primers 31 and 32 are typical primers used in the amplification of a DNA having the partial nucleotide sequence of a raffinose synthase gene. Primer 31 is a sense primer used in the amplification of a DNA having the partial nucleotide sequence of a raffinose synthase gene from mustard and rapeseed and primer 32 is an  
20 antisense primer. Depending upon the purpose, recognition sequences for suitable restriction enzymes can be added to the 5'-termini of these nucleotide sequences.

Primers 33 and 34 are the typical primers used in the amplification of a cDNA of a raffinose synthase gene of mustard.

25 Primers 33 and 34 are both based on the nucleotide sequence of

raffinose synthase gene in the non-translated region. Primer 33 is a sense primer corresponding to the 5'-terminal non-translated region of the mustard-derived raffinose synthase gene. Primer 34 is an antisense primer corresponding to the 3'-terminal non-translated region.

Primers 35 and 36 are typical primers used in the amplification of an open reading frame coding for the amino acid sequence of a raffinose synthase protein in the cDNA of a raffinose synthase gene. Primer 35 is a sense primer corresponding to the 5'-terminus of the above open reading frame. Primer 36 is an antisense primer corresponding to the 3'-terminus. Depending upon the purpose, recognition sequences for suitable restriction enzymes can be added to the 5'-termini of these nucleotide sequences.

List 4:

The nucleotide sequences shown in List 4 are of the primers used in the cloning of a DNA fragment having the present raffinose synthase gene. The base represented by the symbol "i" is inosine. The bases shown in parentheses mean that a mixture of those bases is used in the synthesis. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 5:

The nucleotide sequence shown in List 5 is of the primer used in the analysis of a nucleotide sequence of the present

raffinose synthase gene. 5-SC-2 is used in the analysis of the present nucleotide sequence in the 3'-terminal region.

List 6

5 The nucleotide sequences shown in List 6 are of the primers used in the analysis of the present raffinose synthase gene of beet. The base represented by the symbol "i" is inosine.

10 The bases shown in parentheses mean that a mixture of those bases was used in the synthesis. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 7

15 The nucleotide sequences shown in List 7 are of the primers synthesized on the partial nucleotide sequences of the beet raffinose synthase gene. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 8:

20 The nucleotide sequences shown in List 8 are of the primers used in the analysis of the cDNA nucleotide sequence of a raffinose synthase gene of mustard. The base represented by the symbol "i" is inosine. The bases shown in parentheses mean that a mixture of those bases. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

25

List 9:

The nucleotide sequences shown in List 9 are of the primers synthesized on the partial nucleotide sequences of the mustard raffinose synthase gene. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 10:

The nucleotide sequences shown in List 10 are of the primers used in the analysis of the nucleotide sequences of raffinose synthase gene of mustard and rapeseed. The symbol "RV" as used after the primer number means that the primer referred to by this symbol has an antisense sequence.

List 11:

The nucleotide sequences shown in List 11 are of the primers used in the amplification of 5'-terminal region of a mustard raffinose synthase gene. 11-SacI-BjN is a primer whose SacI restriction site is added to the nucleotide sequence represented by the 4th to 29th nucleotides in SEQ ID NO: 6. 11-SacI-BjINTRV is an antisense primer having a nucleotide sequence corresponding to the nucleotide sequence represented by the 1164th to 1188th nucleotides in SEQ ID NO: 6.

List 12:

The nucleotide sequences shown in List 12 are of the adapters added to a mustard cDNA. These synthetic DNA take a double-stranded form when mixed together because they are complementary strands. This adapter has cohesive ends of cleavage

sites for the restriction enzymes BamHI and SacI on both termini, and contains the restriction sites for the restriction enzymes BamHI and SacI in the double-stranded region.

List 13:

5           The nucleotide sequences shown in List 13 are of the primers used in the confirmation of inserting direction of the mustard-derived raffinose synthase gene. 13-35S-3 is a primer of sense to 35S promoter. 13-B-2RV is an antisense primer having the nucleotide sequence represented by the 593rd to 622nd nucleotides  
10 of SEQ ID NO: 6, 13-B-8 is a sense primer having the nucleotide sequence represented by the 1110th to 1138th nucleotides in SEQ ID NO: 6.

15           As described hereinabove, according to the present invention, it is possible to provide raffinose synthase genes which can be utilized in techniques for varying expression level and activity of raffinose synthase in plants.

SEQUENCE LISTING FREE TEXT

20           SEQ ID NO: 9 to SEQ ID NO: 20: Designed oligonucleotide primer to obtain raffinose synthase gene.

          SEQ ID NO: 21 and SEQ ID NO: 22: Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, r is a or g.

25           SEQ ID NO: 23: Designated oligonucleotide primer to

obtain raffinose synthase gene.

SEQ ID NO: 24 to SEQ ID NO: 27: Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, y is t or c, r is a or g.

5 SEQ ID NO: 28 to SEQ ID NO: 35: Designed oligonucleotide primer to obtain raffinose synthase gene.

SEQ ID NO: 36 and SEQ ID NO: 37: Designed oligonucleotide primer to obtain raffinose synthase gene, n is i, r is a or g.

10 SEQ ID NO: 38 to SEQ ID NO: 48: Designed oligonucleotide primer to obtain raffinose synthase gene.

SEQ ID NO: 49 and SEQ ID NO: 50: Designed oligonucleotide linker to obtain raffinose synthase gene.

15 SEQ ID NO: 51 to SEQ ID NO: 53: Designed oligonucleotide primer to confirm direction of the inserted raffinose synthase gene.

SEQUENCE LISTING

5 <110> Eijiro WATANABE et al.; Sumitomo Chemical Company Limited

<120> Raffinose Synthase Genes and Their Use

<150> JP 10/120550

10 <151> 1998-04-30

<150> JP 10/120551

<151> 1998-04-30

15 <150> JP 10/345590

<151> 1998-12-04

<150> JP 10/351246

<151> 1998-12-10

20 <160> 53

<210> 1

<211> 265

25 <212> PRT

<213> Glycine max

<400> 1

30 Gln Ser Asp His Ala Cys Ala Glu Phe His Ala Ala Ser Arg Ala Ile

5 10 15

Ser Gly Gly Pro Ile Tyr Val Ser Asp Ser Val Gly Lys His Asn Phe

20 25 30

Lys Leu Leu Lys Lys Leu Val Leu Pro Asp Gly Ser Ile Leu Arg Cys

35 40 45

35 Gln His Tyr Ala Leu Pro Thr Arg Asp Cys Leu Phe Val Asp Pro Leu

50 55 60

His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn Leu Asn Lys Cys Ser

65 70 75 80

Gly Val Leu Gly Leu Phe Asn Cys Gln Gly Gly Gly Trp Cys Pro Val

```

25      <210> 2
        <211> 928
        <212> DNA
        <213> Glycine max

30      <220>
        <221> CDS
        <222> (2)... (799)

        <400> 2

35      c caa tct gat cat gct tgt gcc gaa ttc cac gct gct tct aga gcc      46
          Gln Ser Asp His Ala Cys Ala Glu Phe His Ala Ala Ser Arg Ala
                5                10                15
      att tct ggt gga cca att tat gta agc gac tct gtt gga aaa cac aac      94
      Ile Ser Gly Gly Pro Ile Tyr Val Ser Asp Ser Val Gly Lys His Asn

```



		20		25		30		
		ttc aag ttg ctt aag aag ctt gtt cta cct gat ggc tcc att ttg cgg	142					
		Phe Lys Leu Leu Lys Lys Leu Val Leu Pro Asp Gly Ser Ile Leu Arg						
		35		40		45		
5		tgt caa cat tat gca ctt ccc acc cga gac tgc tta ttt gta gat cct	190					
		Cys Gln His Tyr Ala Leu Pro Thr Arg Asp Cys Leu Phe Val Asp Pro						
		50		55		60		
		tta cat gat ggg aaa aca atg ctc aaa att tgg aac ctc aat aaa tgt	238					
		Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp Asn Leu Asn Lys Cys						
10		65		70		75		
		tcc ggg gtt ttg ggt ctg ttc aat tgc caa gga gga ggt tgg tgc cct	286					
		Ser Gly Val Leu Gly Leu Phe Asn Cys Gln Gly Gly Gly Trp Cys Pro						
		80		85		90		95
		ggt act agg cga aac aag agt agc tct gac tat tca cac tcc gtg act	334					
15		Val Thr Arg Arg Asn Lys Ser Ser Ser Asp Tyr Ser His Ser Val Thr						
		100		105		110		
		tgc ttt gca agt cct caa gac att gaa tgg ggc aaa ggg aag cac cca	382					
		Cys Phe Ala Ser Pro Gln Asp Ile Glu Trp Gly Lys Gly Lys His Pro						
		115		120		125		
20		ggt tgc atc aaa ggg gtg gac gta ttt gct gtg tac atg ttt aag gac	430					
		Val Cys Ile Lys Gly Val Asp Val Phe Ala Val Tyr Met Phe Lys Asp						
		130		135		140		
		gac aag ttg aag ctg ctg aag tac aca gag agt gta gaa gtt tct ctt	478					
		Asp Lys Leu Lys Leu Leu Lys Tyr Thr Glu Ser Val Glu Val Ser Leu						
25		145		150		155		
		gag cct ttt agt tgt gag ctt ttg acc gtt tct cca gtg gtg atc tta	526					
		Glu Pro Phe Ser Cys Glu Leu Leu Thr Val Ser Pro Val Val Ile Leu						
		160		165		170		175
		ccc aga aaa tca atc caa ttt gcc cca att gga ttg gta aac atg ctc	574					
30		Pro Arg Lys Ser Ile Gln Phe Ala Pro Ile Gly Leu Val Asn Met Leu						
		180		185		190		
		aac tct ggg ggc tct att atg tca ttg gaa ttt gat caa cag gaa aat	622					
		Asn Ser Gly Gly Ser Ile Met Ser Leu Glu Phe Asp Gln Gln Glu Asn						
		195		200		205		
35		ttg gcg agg att ggg gtg aga gga cat ggg gaa atg agg gta ttt gca	670					
		Leu Ala Arg Ile Gly Val Arg Gly His Gly Glu Met Arg Val Phe Ala						
		210		215		220		
		tca gag aag cca gag agt gtc aag att gat gga gaa tct gtg gaa ttt	718					
		Ser Glu Lys Pro Glu Ser Val Lys Ile Asp Gly Glu Ser Val Glu Phe						

	225		230		235		
	gat tat gtt gat aga acc gtg agg ctc caa gtc tcg tgg cct tgt tct						766
	Asp Tyr Val Asp Arg Thr Val Arg Leu Gln Val Ser Trp Pro Cys Ser						
	240		245		250		255
5	tcg agg ttg tcc gta gtc gag tat ttg ttc tga atcatgattt ggtgtccgag						819
	Ser Arg Leu Ser Val Val Glu Tyr Leu Phe						
			260		265		
	agagccgtgt aatgttcaca taaactgact taagtgcatt aagcaaattcc accttaaata						879
	atagtgcata actttgttcc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa						928
10							
	<210> 3						
	<211> 783						
	<212> PRT						
	<213> Beta vulgaris L.						
15							
	<400> 3						
	Met Ala Pro Ser Phe Ser Lys Glu Asn Ser Lys Thr Cys Asp Glu Val						
		5		10		15	
	Ala Asn His Asp Asp Cys Asn Thr Cys Pro Ile Ile Ser Leu Glu Glu						
20		20		25		30	
	Ser Asn Phe Met Val Asn Gly His Val Ile Leu Ser Gln Val Pro Ser						
		35		40		45	
	Asn Ile Thr Ala Ile Ser Lys Met Gly Phe Asp Gly Leu Phe Val Gly						
		50		55		60	
25	Phe Asp Ala Pro Glu Pro Lys Ala Arg His Val Val Ser Val Gly Gln						
		65		70		75	
	Leu Lys Gly Ile Pro Phe Met Ser Ile Phe Arg Phe Lys Val Trp Trp						
		85		90		95	
	Thr Thr His Trp Thr Gly Ser Asn Gly Arg Asp Leu Glu His Glu Thr						
30		100		105		110	
	Gln Ile Leu Ile Leu Asp Lys Ser Asp Glu Gly Leu Gly Arg Pro Tyr						
		115		120		125	
	Ile Val Ile Leu Pro Leu Ile Glu Gly Pro Phe Arg Ala Ser Leu Gln						
		130		135		140	
35	Pro Gly Ser Val Asp Asp Tyr Val Asp Ile Cys Val Glu Ser Gly Ser						
		145		150		155	
	Thr Lys Val Val Gly Asp Ser Phe Arg Ala Val Leu Tyr Ile Arg Ala						
		165		170		175	
	Gly Pro Asp Pro Phe Lys Leu Ile Lys Asp Thr Met Lys Glu Val Gln						

CCNA 3.0.0.0

35

	Cys	Ala	Tyr	Asn	Ser	Leu	Trp	Met	Gly	Asn	Phe	Ile	His	Pro	Asp	Trp
				500					505					510		
	Asp	Met	Phe	Gln	Ser	Thr	His	Pro	Cys	Ala	Glu	Phe	His	Ala	Ala	Ser
			515					520					525			
5	Arg	Ala	Ile	Ser	Gly	Gly	Pro	Ile	Tyr	Val	Ser	Asp	Val	Val	Gly	Lys
		530					535					540				
	His	Asn	Ile	Pro	Leu	Leu	Lys	Arg	Leu	Val	Leu	Ala	Asp	Gly	Ser	Ile
	545					550					555					560
	Leu	Arg	Cys	Glu	Tyr	His	Ala	Leu	Pro	Thr	Lys	Asp	Cys	Leu	Phe	Val
10					565					570					575	
	Asp	Pro	Leu	His	Asp	Gly	Lys	Thr	Met	Leu	Lys	Ile	Trp	Asn	Leu	Asn
				580					585					590		
	Lys	Tyr	Asn	Gly	Val	Leu	Gly	Val	Phe	Asn	Cys	Gln	Gly	Gly	Gly	Trp
			595					600					605			
15	Ser	Arg	Glu	Ser	Arg	Lys	Asn	Leu	Cys	Phe	Ser	Glu	Tyr	Ser	Lys	Pro
		610					615					620				
	Ile	Ser	Cys	Lys	Thr	Ser	Pro	Lys	Asp	Val	Glu	Trp	Glu	Asn	Gly	His
	625					630					635					640
	Lys	Pro	Phe	Pro	Ile	Lys	Gly	Val	Glu	Cys	Phe	Ala	Met	Tyr	Phe	Thr
20					645					650					655	
	Lys	Glu	Lys	Lys	Leu	Ile	Leu	Ser	Gln	Leu	Ser	Asp	Thr	Ile	Glu	Ile
				660					665					670		
	Ser	Leu	Asp	Pro	Phe	Asp	Tyr	Glu	Leu	Ile	Val	Val	Ser	Pro	Met	Thr
			675					680					685			
25	Ile	Leu	Pro	Trp	Glu	Ser	Ile	Ala	Phe	Ala	Pro	Ile	Gly	Leu	Val	Asn
		690					695					700				
	Met	Leu	Asn	Ala	Gly	Gly	Ala	Val	Lys	Ser	Leu	Asp	Ile	Ser	Glu	Asp
	705					710					715					720
	Asn	Glu	Asp	Lys	Met	Val	Gln	Val	Gly	Ile	Lys	Gly	Ala	Gly	Glu	Met
30					725					730					735	
	Met	Val	Tyr	Ser	Ser	Glu	Lys	Pro	Lys	Ala	Cys	Arg	Val	Asn	Gly	Glu
				740					745					750		
	Asp	Met	Glu	Phe	Glu	Tyr	Glu	Glu	Ser	Met	Ile	Lys	Val	Gln	Val	Thr
			755					760					765			
35	Trp	Asn	His	Asn	Ser	Gly	Gly	Phe	Thr	Thr	Val	Glu	Tyr	Leu	Phe	
		770					775					780			783	

<210> 4

<211> 2690

<212> DNA

<213> Beta vulgaris L.

<220>

5 <221> CDS

<222> (236)... (2587)

<400> 4

	ctaccaaatt ccacaactta aagttcacct caatctttat tccatttttc ctcctaaac	60
10	ttcattgtta agattttgta attgaattca aattcttaat tctgaatttt gtcatttttt	120
	ttgtggggat atttataact atcatattat ttgtgtagat cattctacaa aaaagagagt	180
	gagttttttt agctcttatt tcctaagaaa ttaatagcaa aagttttgca taact atg	238
		Met
	gct cca agc ttt agc aag gaa aat tcc aag acg tgt gat gag gtt gca	286
15	Ala Pro Ser Phe Ser Lys Glu Asn Ser Lys Thr Cys Asp Glu Val Ala	
	5 10 15	
	aac cat gat gat tgc aac acg tgt cca ata att tcc ttg gaa gaa tca	334
	Asn His Asp Asp Cys Asn Thr Cys Pro Ile Ile Ser Leu Glu Glu Ser	
	20 25 30	
20	aac ttc atg gtg aat ggt cac gtg ata ttg tcc caa gtt cca tcc aac	382
	Asn Phe Met Val Asn Gly His Val Ile Leu Ser Gln Val Pro Ser Asn	
	35 40 45	
	atc acg gcc att agt aaa atg ggt ttt gat ggg ctt ttt gtg ggt ttt	430
	Ile Thr Ala Ile Ser Lys Met Gly Phe Asp Gly Leu Phe Val Gly Phe	
25	50 55 60 65	
	gat gct cca gag ccc aag gcc cgg cac gtt gta tcc gtg ggc cag ctc	478
	Asp Ala Pro Glu Pro Lys Ala Arg His Val Val Ser Val Gly Gln Leu	
	70 75 80	
	aag gga att ccc ttc atg agt atc ttc agg ttc aag gta tgg tgg act	526
30	Lys Gly Ile Pro Phe Met Ser Ile Phe Arg Phe Lys Val Trp Trp Thr	
	85 90 95	
	acc cat tgg act ggg tcc aat ggg cgg gac ctt gag cat gag acc caa	574
	Thr His Trp Thr Gly Ser Asn Gly Arg Asp Leu Glu His Glu Thr Gln	
	100 105 110	
35	att ctc atc ctt gat aag tca gat gaa ggt ttg ggc cgt ccc tat att	622
	Ile Leu Ile Leu Asp Lys Ser Asp Glu Gly Leu Gly Arg Pro Tyr Ile	
	115 120 125	
	gtg atc ctc cca ttg atc gaa ggc cca ttt cgg gca tct ctc cag ccg	670
	Val Ile Leu Pro Leu Ile Glu Gly Pro Phe Arg Ala Ser Leu Gln Pro	

	130		135		140		145	
	ggt tct gtt gat gac tat gtg gat ata tgt gtt gag agt ggg tcc act							718
	Gly Ser Val Asp Asp Tyr Val Asp Ile Cys Val Glu Ser Gly Ser Thr							
		150		155		160		
5	aaa gtt gtc gga gac tcg ttc cgg gct gtt ctt tat ata cgg gct ggg							766
	Lys Val Val Gly Asp Ser Phe Arg Ala Val Leu Tyr Ile Arg Ala Gly							
		165		170		175		
	cct gac cca ttt aag tta att aaa gat aca atg aag gaa gtc caa gcc							814
	Pro Asp Pro Phe Lys Leu Ile Lys Asp Thr Met Lys Glu Val Gln Ala							
10		180		185		190		
	cat tta ggg act ttc aaa ctc tta gat gac aaa act cct cca gga ata							862
	His Leu Gly Thr Phe Lys Leu Leu Asp Asp Lys Thr Pro Pro Gly Ile							
		195		200		205		
	gtg gac aag ttt gga tgg tgt aca tgg gat gca ttt tac ctc aaa gta							910
15	Val Asp Lys Phe Gly Trp Cys Thr Trp Asp Ala Phe Tyr Leu Lys Val							
		210		215		220		225
	gag ccw tat ggt gtt tgg gaa gga gtt aaa gga ctc gtc gaa aac ggg							958
	Glu Pro Tyr Gly Val Trp Glu Gly Val Lys Gly Leu Val Glu Asn Gly							
		230		235		240		
20	gtc cca ccc ggt ctc gta ctc att gat gat ggg tgg caa tct att tgt							1006
	Val Pro Pro Gly Leu Val Leu Ile Asp Asp Gly Trp Gln Ser Ile Cys							
		245		250		255		
	cat gac gat gat ccg att acc gac caa gaa ggg ata aac cgg act tct							1054
	His Asp Asp Asp Pro Ile Thr Asp Gln Glu Gly Ile Asn Arg Thr Ser							
25		260		265		270		
	gcc ggc gag caa atg cca tgt aga ttg atc aag tac gag gaa aac ttc							1102
	Ala Gly Glu Gln Met Pro Cys Arg Leu Ile Lys Tyr Glu Glu Asn Phe							
		275		280		285		
	aag ttt agg gac tat aaa agc cca aat att atg ggc cat gag gat cat							1150
30	Lys Phe Arg Asp Tyr Lys Ser Pro Asn Ile Met Gly His Glu Asp His							
		290		295		300		305
	ccc aat atg gga atg agg gcc ttt gtt agg gac ctt aag gag gag ttc							1198
	Pro Asn Met Gly Met Arg Ala Phe Val Arg Asp Leu Lys Glu Glu Phe							
		310		315		320		
35	aaa act gtt gag cat gtg tat gtt tgg cat gct ttt acg ggc tat tgg							1246
	Lys Thr Val Glu His Val Tyr Val Trp His Ala Phe Thr Gly Tyr Trp							
		325		330		335		
	gga ggg gta agg ccc aat gtt cca ggc cta ccr gag gcc caa gta gta							1294
	Gly Gly Val Arg Pro Asn Val Pro Gly Leu Pro Glu Ala Gln Val Val							

		340					345					350						
		acc	cca	aag	ctt	tcc	ccg	ggt	ctt	gag	atg	aca	atg	gaa	gat	cta	gct	1342
		Thr	Pro	Lys	Leu	Ser	Pro	Gly	Leu	Glu	Met	Thr	Met	Glu	Asp	Leu	Ala	
		355					360					365						
5		gtg	gat	aaa	att	gtt	aat	aat	ggt	att	ggg	ctt	gtc	cag	cct	gat	aag	1390
		Val	Asp	Lys	Ile	Val	Asn	Asn	Gly	Ile	Gly	Leu	Val	Gln	Pro	Asp	Lys	
		370					375					380					385	
		gcc	caa	gaa	ctt	tat	gaa	ggg	ttg	cat	tct	cat	ttg	gaa	aat	tgt	ggg	1438
		Ala	Gln	Glu	Leu	Tyr	Glu	Gly	Leu	His	Ser	His	Leu	Glu	Asn	Cys	Gly	
10		390					395					400						
		att	gat	gga	gtc	aaa	gtt	gat	gtc	atc	cat	ttg	ttg	gag	atg	atg	gca	1486
		Ile	Asp	Gly	Val	Lys	Val	Asp	Val	Ile	His	Leu	Leu	Glu	Met	Met	Ala	
		405					410					415						
		gag	gac	tat	gga	gga	aga	gtt	gaa	cta	gca	aaa	aca	tac	tat	aag	gca	1534
15		Glu	Asp	Tyr	Gly	Gly	Arg	Val	Glu	Leu	Ala	Lys	Thr	Tyr	Tyr	Lys	Ala	
		420					425					430						
		ata	aca	gaa	tca	gtg	cgt	aag	cat	ttc	aaa	ggc	aac	ggg	gtg	att	gct	1582
		Ile	Thr	Glu	Ser	Val	Arg	Lys	His	Phe	Lys	Gly	Asn	Gly	Val	Ile	Ala	
		435					440					445						
20		agc	atg	gag	cag	tgc	aac	gat	ttc	atg	ctc	ctt	ggg	act	gag	acc	att	1630
		Ser	Met	Glu	Gln	Cys	Asn	Asp	Phe	Met	Leu	Leu	Gly	Thr	Glu	Thr	Ile	
		450					455					460					465	
		tgt	ctt	ggg	cgc	gtt	ggg	gat	gac	ttt	tgg	cca	act	gat	ccg	tct	gga	1678
		Cys	Leu	Gly	Arg	Val	Gly	Asp	Asp	Phe	Trp	Pro	Thr	Asp	Pro	Ser	Gly	
25		470					475					480						
		gat	ata	aat	ggg	aca	tat	tgg	ctc	caa	ggc	tgt	cat	atg	gtg	cat	tgt	1726
		Asp	Ile	Asn	Gly	Thr	Tyr	Trp	Leu	Gln	Gly	Cys	His	Met	Val	His	Cys	
		485					490					495						
		gcc	tac	aat	agc	tta	tgg	atg	gga	aac	ttt	ata	cac	cct	gac	tgg	gac	1774
30		Ala	Tyr	Asn	Ser	Leu	Trp	Met	Gly	Asn	Phe	Ile	His	Pro	Asp	Trp	Asp	
		500					505					510						
		atg	ttc	caa	tct	aca	cac	cct	tgt	gct	gaa	ttt	cat	gct	gca	tct	cgt	1822
		Met	Phe	Gln	Ser	Thr	His	Pro	Cys	Ala	Glu	Phe	His	Ala	Ala	Ser	Arg	
		515					520					525						
35		gcg	att	tct	ggg	gga	cca	att	tat	gtt	agt	gat	gtt	gtt	ggc	aag	cat	1870
		Ala	Ile	Ser	Gly	Gly	Pro	Ile	Tyr	Val	Ser	Asp	Val	Val	Gly	Lys	His	
		530					535					540					545	
		aac	atc	ccc	ttg	ctc	aaa	agg	ctc	gtc	ttg	gct	gat	ggg	tcg	atc	ctt	1918
		Asn	Ile	Pro	Leu	Leu	Lys	Arg	Leu	Val	Leu	Ala	Asp	Gly	Ser	Ile	Leu	

				550					555					560					
		cgt	tgc	gag	tac	cat	gca	ctt	cct	act	aag	gat	tgc	cta	ttt	gta	gat		1966
		Arg	Cys	Glu	Tyr	His	Ala	Leu	Pro	Thr	Lys	Asp	Cys	Leu	Phe	Val	Asp		
				565						570					575				
5		cct	ttg	cac	gat	ggc	aaa	aca	atg	ctc	aaa	att	tgg	aac	ctc	aac	aag		2014
		Pro	Leu	His	Asp	Gly	Lys	Thr	Met	Leu	Lys	Ile	Trp	Asn	Leu	Asn	Lys		
				580						585					590				
		tac	aat	gga	gtg	ctt	gga	gtc	ttc	aat	tgc	caa	gga	gga	ggg	tgg	agc		2062
		Tyr	Asn	Gly	Val	Leu	Gly	Val	Phe	Asn	Cys	Gln	Gly	Gly	Gly	Trp	Ser		
10			595					600					605						
		cgt	gag	tct	cga	aaa	aat	cta	tgt	ttc	tca	gag	tat	tca	aaa	cct	att		2110
		Arg	Glu	Ser	Arg	Lys	Asn	Leu	Cys	Phe	Ser	Glu	Tyr	Ser	Lys	Pro	Ile		
		610				615					620				625				
		tcc	tgc	aag	aca	agt	cca	aaa	gat	gtt	gaa	tgg	gag	aac	gga	cac	aag		2158
15		Ser	Cys	Lys	Thr	Ser	Pro	Lys	Asp	Val	Glu	Trp	Glu	Asn	Gly	His	Lys		
				630						635					640				
		cca	ttc	ccc	atc	aaa	gga	gtg	gaa	tgt	ttt	gcc	atg	tac	ttc	acc	aag		2206
		Pro	Phe	Pro	Ile	Lys	Gly	Val	Glu	Cys	Phe	Ala	Met	Tyr	Phe	Thr	Lys		
				645						650					655				
20		gaa	aaa	aag	cta	atc	ctc	tca	caa	cta	tct	gac	acc	att	gaa	ata	tca		2254
		Glu	Lys	Lys	Leu	Ile	Leu	Ser	Gln	Leu	Ser	Asp	Thr	Ile	Glu	Ile	Ser		
			660					665					670						
		ctt	gat	ccc	ttc	gat	tac	gag	ctt	att	gta	gtc	tct	ccg	atg	aca	att		2302
		Leu	Asp	Pro	Phe	Asp	Tyr	Glu	Leu	Ile	Val	Val	Ser	Pro	Met	Thr	Ile		
25			675					680					685						
		cta	ccc	tgg	gag	tcg	atc	gca	ttt	gca	ccc	ata	gga	tta	gta	aac	atg		2350
		Leu	Pro	Trp	Glu	Ser	Ile	Ala	Phe	Ala	Pro	Ile	Gly	Leu	Val	Asn	Met		
		690				695					700				705				
		ctc	aac	gcc	gga	ggg	gca	gtc	aag	tct	ttg	gac	atc	agt	gag	gat	aat		2398
30		Leu	Asn	Ala	Gly	Gly	Ala	Val	Lys	Ser	Leu	Asp	Ile	Ser	Glu	Asp	Asn		
				710						715					720				
		gag	gat	aag	atg	gtt	cag	gtt	ggt	att	aaa	ggg	gcc	gga	gaa	atg	atg		2446
		Glu	Asp	Lys	Met	Val	Gln	Val	Gly	Ile	Lys	Gly	Ala	Gly	Glu	Met	Met		
				725						730					735				
35		gtt	tat	tca	tca	gaa	aag	cca	aaa	gcg	tgt	aga	gtt	aat	gga	gaa	gac		2494
		Val	Tyr	Ser	Ser	Glu	Lys	Pro	Lys	Ala	Cys	Arg	Val	Asn	Gly	Glu	Asp		
			740					745					750						
		atg	gag	ttt	gag	tat	gaa	gag	agc	atg	att	aag	gtt	caa	gtt	aca	tgg		2542
		Met	Glu	Phe	Glu	Tyr	Glu	Glu	Ser	Met	Ile	Lys	Val	Gln	Val	Thr	Trp		



	755		760		765		
	aac cat aac tca ggt ggt ttt acc act gtt gag tac tta ttt tga gcttg					2592	
	Asn His Asn Ser Gly Gly Phe Thr Thr Val Glu Tyr Leu Phe						
	770		775		780		
5	aagctaattct aagtcctttac ttaatgagtg atgtaactga gtagttgact tgagagtaca					2652	
	gtatgtgtga agcttattat tccaaaaaaa aaaaaaaa					2690	
	<210> 5						
	<211> 777						
10	<212> PRT						
	<213> Brassica juncea						
	<400> 5						
15	Met Ala Pro Pro Ser Val Ile Lys Ser Asp Ala Ala Val Asn Gly Ile						
		5		10		15	
	Asp Leu Ser Gly Lys Pro Leu Phe Arg Leu Glu Gly Ser Asp Leu Leu						
		20		25		30	
	Ala Asn Gly His Val Val Leu Thr Asp Val Pro Val Asn Val Thr Val						
		35		40		45	
20	Thr Ala Ser Pro Tyr Leu Ala Asp Lys Asp Gly Glu Pro Val Asp Ala						
		50		55		60	
	Ser Ala Gly Ser Phe Ile Gly Phe Asn Leu Asp Gly Glu Pro Arg Ser						
		65		70		75	
	Arg His Val Ala Ser Ile Gly Lys Leu Arg Asp Ile Arg Phe Met Ser						
25		85		90		95	
	Ile Phe Arg Phe Lys Val Trp Trp Thr Thr His Trp Val Gly Ser Lys						
		100		105		110	
	Gly Ser Asp Ile Glu Asn Glu Thr Gln Ile Ile Ile Leu Glu Asn Ser						
		115		120		125	
30	Gly Ser Gly Arg Pro Tyr Val Leu Leu Leu Pro Leu Leu Glu Gly Ser						
		130		135		140	
	Phe Arg Ser Ser Phe Gln Pro Gly Glu Asp Asp Asp Val Ala Val Cys						
		145		150		155	
	Val Glu Ser Gly Ser Thr Gln Val Thr Gly Ser Glu Phe Arg Gln Val						
35		165		170		175	
	Val Tyr Val His Ala Gly Asp Asp Pro Phe Lys Leu Val Lys Asp Ala						
		180		185		190	
	Met Lys Val Val Arg Val His Met Asn Thr Phe Lys Leu Leu Glu Glu						
		195		200		205	

	Lys	Thr	Pro	Pro	Gly	Ile	Val	Asp	Lys	Phe	Gly	Trp	Cys	Thr	Trp	Asp
	210						215					220				
	Ala	Phe	Tyr	Leu	Thr	Val	Asn	Pro	Asp	Gly	Val	His	Lys	Gly	Val	Lys
	225					230				235						240
5	Cys	Leu	Val	Asp	Gly	Gly	Cys	Pro	Pro	Gly	Leu	Val	Leu	Ile	Asp	Asp
					245					250					255	
	Gly	Trp	Gln	Ser	Ile	Gly	His	Asp	Ser	Asp	Gly	Ile	Asp	Val	Glu	Gly
				260					265					270		
10	Met	Ser	Cys	Thr	Val	Ala	Gly	Glu	Gln	Met	Pro	Cys	Arg	Leu	Leu	Lys
			275					280					285			
	Phe	Gln	Glu	Asn	Phe	Lys	Phe	Arg	Asp	Tyr	Val	Ser	Pro	Lys	Asp	Lys
	290						295					300				
	Asn	Glu	Val	Gly	Met	Lys	Ala	Phe	Val	Arg	Asp	Leu	Lys	Glu	Glu	Phe
	305					310					315					320
15	Ser	Thr	Val	Asp	Tyr	Ile	Tyr	Val	Trp	His	Ala	Leu	Cys	Gly	Tyr	Trp
					325					330					335	
	Gly	Gly	Leu	Arg	Pro	Gly	Ala	Pro	Thr	Leu	Pro	Pro	Ser	Thr	Ile	Val
				340					345					350		
20	Arg	Pro	Glu	Leu	Ser	Pro	Gly	Leu	Lys	Leu	Thr	Met	Gln	Asp	Leu	Ala
			355					360					365			
	Val	Asp	Lys	Ile	Val	Asp	Thr	Gly	Ile	Gly	Phe	Val	Ser	Pro	Asp	Met
	370						375					380				
	Ala	Asn	Glu	Phe	Tyr	Glu	Gly	Leu	His	Ser	His	Leu	Gln	Asn	Val	Gly
	385					390					395					400
25	Ile	Asp	Gly	Val	Lys	Val	Asp	Val	Ile	His	Ile	Leu	Glu	Met	Leu	Cys
					405					410					415	
	Glu	Lys	Tyr	Gly	Gly	Arg	Val	Asp	Leu	Ala	Lys	Ala	Tyr	Phe	Lys	Ala
				420					425					430		
30	Leu	Thr	Ser	Ser	Val	Asn	Lys	His	Phe	Asp	Gly	Asn	Gly	Val	Ile	Ala
			435					440					445			
	Ser	Met	Glu	His	Cys	Asn	Asp	Phe	Met	Phe	Leu	Gly	Thr	Glu	Ala	Ile
	450						455					460				
	Ser	Leu	Gly	Arg	Val	Gly	Asp	Asp	Phe	Trp	Cys	Thr	Asp	Pro	Ser	Gly
	465					470					475					480
35	Asp	Ile	Asn	Gly	Thr	Tyr	Trp	Leu	Gln	Gly	Cys	His	Met	Val	His	Cys
					485					490					495	
	Ala	Tyr	Asn	Ser	Leu	Trp	Met	Gly	Asn	Phe	Ile	Gln	Pro	Asp	Trp	Asp
			500						505					510		
	Met	Phe	Gln	Ser	Thr	His	Pro	Cys	Ala	Glu	Phe	His	Ala	Ala	Ser	Arg

35

<220>

<221> CDS

<222> (134)... (2467)

5 <400> 6

accaatccaa aatctcatca aataatcgca attaggggaa gtttacaaga ttcacatcatct 60  
ccgttactat ataactacgc tcttcttctt tgcctaatc caacttaacc taaaaaccac 120  
tctatcagcg aaa atg gct cca ccg agc gta att aaa tcc gat gct gca 169  
Met Ala Pro Pro Ser Val Ile Lys Ser Asp Ala Ala

10 5 10  
gtc aac ggc att gac ctc tcc gga aag ccg ctt ttc cgg cta gag ggt 217  
Val Asn Gly Ile Asp Leu Ser Gly Lys Pro Leu Phe Arg Leu Glu Gly  
15 20 25  
tcc gat ctc cta gcc aat ggt cac gtt gtc tta acc gat gta ccg gtt 265  
Ser Asp Leu Leu Ala Asn Gly His Val Val Leu Thr Asp Val Pro Val  
30 35 40  
aac gtg act gtc act gct tca cct tac cta gct gac aaa gac gga gaa 313  
Asn Val Thr Val Thr Ala Ser Pro Tyr Leu Ala Asp Lys Asp Gly Glu  
45 50 55 60  
20 ccg gtt gac gcc tcc gct ggt tca ttc atc ggg ttt aat ctc gac ggt 361  
Pro Val Asp Ala Ser Ala Gly Ser Phe Ile Gly Phe Asn Leu Asp Gly  
65 70 75  
gag cca cga agc cgc cac gtg gcg tcc atc ggt aaa ctc agg gat att 409  
Glu Pro Arg Ser Arg His Val Ala Ser Ile Gly Lys Leu Arg Asp Ile  
80 85 90  
25 cga ttc atg agc ata ttc cgt ttc aag gtt tgg tgg act act cac tgg 457  
Arg Phe Met Ser Ile Phe Arg Phe Lys Val Trp Trp Thr Thr His Trp  
95 100 105  
gtc ggt tcc aaa gga tcc gac atc gag aac gag acc cag atc atc atc 505  
30 Val Gly Ser Lys Gly Ser Asp Ile Glu Asn Glu Thr Gln Ile Ile Ile  
110 115 120  
ctc gag aac tcc ggg tcg ggt cgt cct tat gtt ctt ctt ctg ccg ctt 553  
Leu Glu Asn Ser Gly Ser Gly Arg Pro Tyr Val Leu Leu Leu Pro Leu  
125 130 135 140  
35 ctt gaa ggc tct ttc cgt tca tcc ttt cag cct ggg gaa gac gat gac 601  
Leu Glu Gly Ser Phe Arg Ser Ser Phe Gln Pro Gly Glu Asp Asp Asp  
145 150 155  
gtg gcg gtt tgt gtc gaa tcc ggg tcg acc cag gtg acc ggg tcg gag 649  
Val Ala Val Cys Val Glu Ser Gly Ser Thr Gln Val Thr Gly Ser Glu

CCDS:CCDS33266.1

		160		165		170		
		ttt cgt caa gtt gtg tat gtt cac gcc gga gac gat ccg ttc aag ctc						697
		Phe Arg Gln Val Val Tyr Val His Ala Gly Asp Asp Pro Phe Lys Leu						
		175		180		185		
5		gtg aaa gac gcg atg aag gtg gtt agg gtt cat atg aac acc ttc aag						745
		Val Lys Asp Ala Met Lys Val Val Arg Val His Met Asn Thr Phe Lys						
		190		195		200		
		ctc ttg gaa gag aag acr ccg ccg gga atc gtc gat aag ttc ggg tgg						793
		Leu Leu Glu Glu Lys Thr Pro Pro Gly Ile Val Asp Lys Phe Gly Trp						
10		205		210		215		220
		tgc acg tgg gat gcg ttt tat ttg acg gtg aac cct gac gga gtt cat						841
		Cys Thr Trp Asp Ala Phe Tyr Leu Thr Val Asn Pro Asp Gly Val His						
		225		230		235		
		aag ggt gtt aag tgt ctc gtc gac ggt ggt tgt ccg ccg gga ttg gtc						889
15		Lys Gly Val Lys Cys Leu Val Asp Gly Gly Cys Pro Pro Gly Leu Val						
		240		245		250		
		cta atc gac gac ggt tgg caa tcg att gga cat gac tcc gat ggt atc						937
		Leu Ile Asp Asp Gly Trp Gln Ser Ile Gly His Asp Ser Asp Gly Ile						
		255		260		265		
20		gat gtt gaa ggg atg agt tgt acc gtc gcc ggg gag caa atg cct tgc						985
		Asp Val Glu Gly Met Ser Cys Thr Val Ala Gly Glu Gln Met Pro Cys						
		270		275		280		
		agg ctt ctg aaa ttt caa gag aac ttc aag ttc aga gac tac gtc tct						1033
		Arg Leu Leu Lys Phe Gln Glu Asn Phe Lys Phe Arg Asp Tyr Val Ser						
25		285		290		295		300
		ccg aaa gac aaa aac gaa gtc ggg atg aaa gct ttc gtc aga gat ctg						1081
		Pro Lys Asp Lys Asn Glu Val Gly Met Lys Ala Phe Val Arg Asp Leu						
		305		310		315		
		aaa gaa gaa ttc tcc acc gtt gat tac atc tac gtc tgg cac gcg ctt						1129
30		Lys Glu Glu Phe Ser Thr Val Asp Tyr Ile Tyr Val Trp His Ala Leu						
		320		325		330		
		tgc ggc tac tgg ggt ggt ctt cgt ccc gga gct cct act ctt ccg ccc						1177
		Cys Gly Tyr Trp Gly Gly Leu Arg Pro Gly Ala Pro Thr Leu Pro Pro						
		335		340		345		
35		tca act att gtc cgg cca gag ctc tcg ccg ggg ctt aag ttg acg atg						1225
		Ser Thr Ile Val Arg Pro Glu Leu Ser Pro Gly Leu Lys Leu Thr Met						
		350		355		360		
		caa gat ctc gcc gtt gat aag att gtc gat acc gga atc gga ttc gtc						1273
		Gln Asp Leu Ala Val Asp Lys Ile Val Asp Thr Gly Ile Gly Phe Val						

	365		370		375		380		
	tcg ccg gac atg gcg aat gag ttt tac gaa ggt ctt cac tct cat ctt								1321
	Ser Pro Asp Met Ala Asn Glu Phe Tyr Glu Gly Leu His Ser His Leu								
		385		390		395			
5	caa aac gtc ggt att gac ggc gtt aaa gtt gac gtc atc cac ata ttg								1369
	Gln Asn Val Gly Ile Asp Gly Val Lys Val Asp Val Ile His Ile Leu								
		400		405		410			
	gag atg ttg tgc gag aaa tat ggc ggg aga gta gac ttg gct aaa gct								1417
	Glu Met Leu Cys Glu Lys Tyr Gly Gly Arg Val Asp Leu Ala Lys Ala								
10		415		420		425			
	tac ttc aag gcg tta act tcc tca gtg aat aag cat ttt gac ggt aac								1465
	Tyr Phe Lys Ala Leu Thr Ser Ser Val Asn Lys His Phe Asp Gly Asn								
		430		435		440			
	ggc gtt atc gct agc atg gag cac tgt aat gat ttc atg ttc ctt gga								1513
15	Gly Val Ile Ala Ser Met Glu His Cys Asn Asp Phe Met Phe Leu Gly								
		445		450		455		460	
	acc gaa gcc atc tct cta ggt cgt gtc ggt gat gac ttt tgg tgc acg								1561
	Thr Glu Ala Ile Ser Leu Gly Arg Val Gly Asp Asp Phe Trp Cys Thr								
		465		470		475			
20	gat cca tca ggc gac ata aac ggc aca tat tgg ctg caa gga tgc cac								1609
	Asp Pro Ser Gly Asp Ile Asn Gly Thr Tyr Trp Leu Gln Gly Cys His								
		480		485		490			
	atg gtc cac tgt gcc tac aac agt ctt tgg atg gga aat ttc atc cag								1657
	Met Val His Cys Ala Tyr Asn Ser Leu Trp Met Gly Asn Phe Ile Gln								
25		495		500		505			
	cct gat tgg gac atg ttt cag tcc aca cat cct tgt gct gag ttc cat								1705
	Pro Asp Trp Asp Met Phe Gln Ser Thr His Pro Cys Ala Glu Phe His								
		510		515		520			
	gct gct tct cgt gcc atc tcc ggt ggg ccc att tac atc agc gat tgt								1753
30	Ala Ala Ser Arg Ala Ile Ser Gly Gly Pro Ile Tyr Ile Ser Asp Cys								
		525		530		535		540	
	gtg ggc cag cac gat ttc gat ctc ttg aag cga ctc gtc ttg cct gac								1801
	Val Gly Gln His Asp Phe Asp Leu Leu Lys Arg Leu Val Leu Pro Asp								
		545		550		555			
35	ggt tcg att ttg agg tgt gag cac tat gca ctc cca act cgt gac cgt								1849
	Gly Ser Ile Leu Arg Cys Glu His Tyr Ala Leu Pro Thr Arg Asp Arg								
		560		565		570			
	ctc ttt gaa gac cct ctt cat gat ggc aaa acc atg ctc aag att tgg								1897
	Leu Phe Glu Asp Pro Leu His Asp Gly Lys Thr Met Leu Lys Ile Trp								

		575					580					585						
		aac	ttg	aac	aag	tac	act	gga	att	att	gga	gca	ttc	aac	tgc	caa	gga	1945
		Asn	Leu	Asn	Lys	Tyr	Thr	Gly	Ile	Ile	Gly	Ala	Phe	Asn	Cys	Gln	Gly	
		590						595				600						
5		gga	gga	tgg	tgc	aga	gaa	acc	cga	cgc	aac	caa	tgc	ttc	tcc	caa	tgc	1993
		Gly	Gly	Trp	Cys	Arg	Glu	Thr	Arg	Arg	Asn	Gln	Cys	Phe	Ser	Gln	Cys	
		605					610				615					620		
		ggt	aac	acg	tta	acc	gcc	aca	aca	aat	cct	aag	gac	ggt	gaa	tgg	aac	2041
		Val	Asn	Thr	Leu	Thr	Ala	Thr	Thr	Asn	Pro	Lys	Asp	Val	Glu	Trp	Asn	
10						625					630					635		
		agt	ggg	aac	aac	cca	atc	tcc	ggt	gaa	aac	ggt	gaa	gag	ttt	gct	ttg	2089
		Ser	Gly	Asn	Asn	Pro	Ile	Ser	Val	Glu	Asn	Val	Glu	Glu	Phe	Ala	Leu	
						640					645				650			
		ttc	ttg	tct	cag	tct	aag	aag	ctt	gtg	ttg	tct	gga	cca	aac	gat	gat	2137
15		Phe	Leu	Ser	Gln	Ser	Lys	Lys	Leu	Val	Leu	Ser	Gly	Pro	Asn	Asp	Asp	
						655					660				665			
		ctc	gag	atc	act	ttg	gag	cct	ttc	aag	ttt	gag	cta	atc	act	gtc	tca	2185
		Leu	Glu	Ile	Thr	Leu	Glu	Pro	Phe	Lys	Phe	Glu	Leu	Ile	Thr	Val	Ser	
						670							680					
20		cca	ggt	gtc	act	att	gag	ggt	agt	tgc	ggt	cag	ttt	gct	cca	atc	gga	2233
		Pro	Val	Val	Thr	Ile	Glu	Gly	Ser	Ser	Val	Gln	Phe	Ala	Pro	Ile	Gly	
		685					690					695				700		
		ttg	ggt	aac	atg	cta	aac	act	agc	ggt	gca	att	cga	tcc	ttg	gtg	tat	2281
		Leu	Val	Asn	Met	Leu	Asn	Thr	Ser	Gly	Ala	Ile	Arg	Ser	Leu	Val	Tyr	
25						705					710					715		
		cat	gag	gaa	tcc	ggt	gag	att	gga	ggt	cgt	ggt	gct	gga	gag	ttc	agg	2329
		His	Glu	Glu	Ser	Val	Glu	Ile	Gly	Val	Arg	Gly	Ala	Gly	Glu	Phe	Arg	
						720					725				730			
		ggt	tat	gca	tca	agg	aaa	cct	gcg	agc	tgc	aaa	att	gat	ggt	gaa	ggt	2377
30		Val	Tyr	Ala	Ser	Arg	Lys	Pro	Ala	Ser	Cys	Lys	Ile	Asp	Gly	Glu	Val	
						735					740				745			
		ggt	gag	ttt	gga	tac	gaa	gag	tca	atg	gtg	atg	ggt	caa	gtg	cct	tgg	2425
		Val	Glu	Phe	Gly	Tyr	Glu	Glu	Ser	Met	Val	Met	Val	Gln	Val	Pro	Trp	
						750					755				760			
35		tct	gca	ccc	gag	ggt	ttg	tct	tct	att	aag	tat	gag	ttt	tag	agttttccga	2476	
		Ser	Ala	Pro	Glu	Gly	Leu	Ser	Ser	Ile	Lys	Tyr	Glu	Phe				

taagtgattt gtttttgaac tgttttttagt ggtgtaattt gtactgcccc tattattttt 2656  
catattttatt tgtgaaagat aaaaaaaaaa aaaa 2690

<210> 7

5 <211> 572

<212> PRT

<213> Brassica napus

<400> 7

10 Leu Glu Glu Lys Thr Pro Pro Gly Ile Val Asp Lys Phe Gly Trp Cys  
5 10 15  
Thr Trp Asp Ala Phe Tyr Leu Thr Val Asn Pro Asp Gly Val His Lys  
20 25 30  
Gly Val Lys Cys Leu Val Asp Gly Gly Cys Pro Pro Gly Leu Val Leu  
15 35 40 45  
Ile Asp Asp Gly Trp Gln Ser Ile Gly His Asp Ser Asp Gly Ile Asp  
50 55 60  
Val Glu Gly Met Ser Cys Thr Val Ala Gly Glu Gln Met Pro Cys Arg  
65 70 75 80  
20 Leu Pro Lys Phe Gln Glu Asn Phe Lys Phe Arg Asp Tyr Val Ser Pro  
85 90 95  
Lys Asp Lys Asn Glu Val Gly Met Lys Ala Phe Val Arg Asp Leu Lys  
100 105 110  
Glu Glu Phe Ser Thr Val Asp Tyr Ile Tyr Val Trp His Ala Leu Cys  
25 115 120 125  
Gly Tyr Trp Gly Gly Leu Arg Pro Gly Ala Pro Thr Leu Pro Pro Ser  
130 135 140  
Thr Ile Val Arg Pro Glu Leu Ser Pro Gly Leu Lys Leu Thr Met Gln  
145 150 155 160  
30 Asp Leu Ala Val Asp Lys Ile Ile Asp Thr Gly Ile Gly Phe Val Ser  
165 170 175  
Pro Asp Met Ala Asn Glu Phe Tyr Glu Gly Leu His Ser His Leu Gln  
180 185 190  
Asn Val Gly Ile Asn Gly Val Lys Val Asp Val Ile His Ile Leu Glu  
35 195 200 205  
Met Leu Cys Glu Lys Tyr Gly Gly Arg Val Asp Leu Ala Lys Ala Tyr  
210 215 220  
Phe Lys Ala Leu Thr Ser Ser Val Asn Lys His Phe Asp Gly Asn Ala  
225 230 235 240

CCGAGGCG



	Val	Ile	Ala	Ser	Met	Glu	His	Cys	Asn	Asp	Phe	Met	Phe	Leu	Gly	Thr
					245					250					255	
	Glu	Ala	Ile	Ser	Leu	Gly	Arg	Val	Gly	Asp	Asp	Phe	Trp	Cys	Thr	Asp
				260					265					270		
5	Pro	Ser	Gly	Asp	Ile	Asn	Gly	Thr	Tyr	Trp	Leu	Gln	Gly	Cys	His	Met
			275					280					285			
	Val	His	Cys	Ala	Tyr	Asn	Ser	Leu	Trp	Met	Gly	Asn	Phe	Ile	Gln	Pro
	290						295					300				
10	Asp	Trp	Asp	Met	Phe	Gln	Ser	Thr	His	Pro	Cys	Ala	Glu	Phe	His	Ala
	305					310					315					320
	Ala	Ser	Arg	Ala	Ile	Ser	Gly	Gly	Pro	Ile	Tyr	Ile	Ser	Asp	Cys	Val
					325					330					335	
	Gly	Gln	His	Asp	Phe	Asp	Leu	Leu	Arg	Arg	Leu	Val	Leu	Pro	Asp	Gly
				340					345					350		
15	Ser	Ile	Leu	Arg	Cys	Glu	Tyr	Tyr	Ala	Leu	Pro	Thr	Arg	Asp	Arg	Leu
			355					360					365			
	Phe	Glu	Asp	Pro	Leu	His	Asp	Gly	Lys	Thr	Met	Leu	Lys	Ile	Trp	Asn
	370						375					380				
20	Leu	Asn	Lys	Tyr	Thr	Gly	Ile	Ile	Gly	Ala	Phe	Asn	Cys	Gln	Gly	Gly
	385					390					395					400
	Gly	Trp	Cys	Arg	Glu	Thr	Arg	Arg	Asp	Gln	Cys	Phe	Ser	Gln	Cys	Val
				405						410					415	
	Asn	Thr	Leu	Thr	Ala	Thr	Thr	Asn	Pro	Asn	Asp	Val	Glu	Trp	Asn	Ser
				420					425					430		
25	Gly	Asn	Asn	Pro	Ile	Ser	Ile	Glu	Asn	Val	Glu	Glu	Phe	Ala	Leu	Phe
		435						440					445			
	Leu	Ser	Gln	Ser	Lys	Lys	Leu	Val	Leu	Ser	Gly	Gln	Asn	Asp	Asp	Leu
	450						455					460				
30	Glu	Ile	Thr	Leu	Glu	Pro	Phe	Lys	Phe	Glu	Leu	Ile	Thr	Val	Ser	Pro
	465					470					475					480
	Val	Val	Thr	Ile	Glu	Gly	Ser	Ser	Val	Gln	Phe	Ala	Pro	Ile	Gly	Leu
				485						490					495	
	Val	Asn	Met	Leu	Asn	Thr	Ser	Gly	Ala	Ile	Arg	Ser	Leu	Val	Tyr	His
			500						505					510		
35	Glu	Glu	Ser	Val	Glu	Ile	Gly	Val	Arg	Gly	Ala	Gly	Glu	Phe	Arg	Val
			515					520					525			
	Tyr	Ala	Ser	Lys	Lys	Pro	Val	Ser	Cys	Lys	Ile	Asp	Gly	Glu	Asp	Val
	530						535					540				
	Glu	Phe	Gly	Tyr	Glu	Glu	Ser	Met	Val	Met	Val	Gln	Val	Pro	Trp	Ser

<210> 8  
<211> 1762  
<212> DNA  
<213> Brassica napus

<220>  
<221> CDS  
<222> (1)... (1719)

<400> 8

ttg gaa gaa aaa acg ccg ccg gga atc gtc gat aag ttc ggg tgg tgc	48
Leu Glu Glu Lys Thr Pro Pro Gly Ile Val Asp Lys Phe Gly Trp Cys	
5 10 15	
acg tgg gat gcg ttt tat ttg acg gtg aac cct gac gga gtt cat aag	96
Thr Trp Asp Ala Phe Tyr Leu Thr Val Asn Pro Asp Gly Val His Lys	
20 25 30	
ggg gtt aag tgt ctc gtc gac ggt ggt tgt ccg ccg gga ttg gtc cta	144
Gly Val Lys Cys Leu Val Asp Gly Gly Cys Pro Pro Gly Leu Val Leu	
35 40 45	
atc gac gac ggt tgg caa tcg att gga cat gac tcc gat ggt atc gat	192
Ile Asp Asp Gly Trp Gln Ser Ile Gly His Asp Ser Asp Gly Ile Asp	
50 55 60	
gtt gaa ggg atg agt tgt acc gtc gcc ggg gag caa atg cct tgc agg	240
Val Glu Gly Met Ser Cys Thr Val Ala Gly Glu Gln Met Pro Cys Arg	
65 70 75 80	
ctt ccg aaa ttt caa gag aac ttc aag ttc aga gac tac gtc tct ccg	288
Leu Pro Lys Phe Gln Glu Asn Phe Lys Phe Arg Asp Tyr Val Ser Pro	
85 90 95	
aaa gac aaa aac gaa gtc ggg atg aaa gct ttc gtc aga gat ctg aaa	336
Lys Asp Lys Asn Glu Val Gly Met Lys Ala Phe Val Arg Asp Leu Lys	
100 105 110	
gaa gaa ttc tcc acc gtt gat tac atc tac gtc tgg cac gcg ctt tgc	384
Glu Glu Phe Ser Thr Val Asp Tyr Ile Tyr Val Trp His Ala Leu Cys	
115 120 125	
ggg tac tgg ggw ggt ctt cgt ccc gga gct cct act ctt ccg ccs tcr	432

	Gly	Tyr	Trp	Gly	Gly	Leu	Arg	Pro	Gly	Ala	Pro	Thr	Leu	Pro	Pro	Ser	
	130						135					140					
	act	att	gtc	cgr	cca	gag	ctc	tgc	cgc	ggg	ctt	aag	ttg	acg	atg	caa	480
	Thr	Ile	Val	Arg	Pro	Glu	Leu	Ser	Pro	Gly	Leu	Lys	Leu	Thr	Met	Gln	
5	145					150					155					160	
	gat	ctc	gcc	gtt	gat	aag	atc	atc	gat	acc	gga	atc	gga	ttc	gtc	tgc	528
	Asp	Leu	Ala	Val	Asp	Lys	Ile	Ile	Asp	Thr	Gly	Ile	Gly	Phe	Val	Ser	
					165					170					175		
	ccg	gac	atg	gcg	aac	gag	ttt	tac	gaa	ggg	ctt	cac	tct	cat	ctt	caa	576
10	Pro	Asp	Met	Ala	Asn	Glu	Phe	Tyr	Glu	Gly	Leu	His	Ser	His	Leu	Gln	
				180					185					190			
	aac	gtc	ggc	att	aac	ggc	gtt	aaa	gtt	gac	gtt	atc	cac	ata	ctg	gag	624
	Asn	Val	Gly	Ile	Asn	Gly	Val	Lys	Val	Asp	Val	Ile	His	Ile	Leu	Glu	
			195					200				205					
15	atg	ttg	tgc	gag	aaa	tat	ggc	ggg	aga	gtt	gac	ttg	gct	aaa	gct	tac	672
	Met	Leu	Cys	Glu	Lys	Tyr	Gly	Gly	Arg	Val	Asp	Leu	Ala	Lys	Ala	Tyr	
		210					215				220						
	ttc	aag	gcg	tta	acg	tgc	tca	gtg	aat	aag	cat	ttt	gac	ggc	aac	gcc	720
	Phe	Lys	Ala	Leu	Thr	Ser	Ser	Val	Asn	Lys	His	Phe	Asp	Gly	Asn	Ala	
20	225					230					235					240	
	gtt	atc	gcc	agc	atg	gag	cac	tgt	aat	gac	ttc	atg	ttc	ctt	gga	acc	768
	Val	Ile	Ala	Ser	Met	Glu	His	Cys	Asn	Asp	Phe	Met	Phe	Leu	Gly	Thr	
					245					250					255		
	gaa	gcc	atc	tct	cta	ggg	cgt	gtc	ggg	gat	gac	ttt	tgg	tgc	acg	gat	816
25	Glu	Ala	Ile	Ser	Leu	Gly	Arg	Val	Gly	Asp	Asp	Phe	Trp	Cys	Thr	Asp	
			260						265				270				
	cca	tct	ggc	gac	att	aac	ggc	acg	tat	tgg	ctg	caa	gga	tgt	cac	atg	864
	Pro	Ser	Gly	Asp	Ile	Asn	Gly	Thr	Tyr	Trp	Leu	Gln	Gly	Cys	His	Met	
			275					280				285					
30	gtc	cac	tgt	gcc	tac	aac	agt	ctt	tgg	atg	gga	aat	ttc	atc	cag	cct	912
	Val	His	Cys	Ala	Tyr	Asn	Ser	Leu	Trp	Met	Gly	Asn	Phe	Ile	Gln	Pro	
		290					295				300						
	gat	tgg	gac	atg	ttt	cag	tcc	aca	cat	cct	tgt	gct	gag	ttc	cat	gct	960
	Asp	Trp	Asp	Met	Phe	Gln	Ser	Thr	His	Pro	Cys	Ala	Glu	Phe	His	Ala	
35	305					310					315					320	
	gct	tca	cgt	gcc	atc	tcc	ggg	ggg	ccc	att	tac	atc	agc	gat	tgt	gtg	1008
	Ala	Ser	Arg	Ala	Ile	Ser	Gly	Gly	Pro	Ile	Tyr	Ile	Ser	Asp	Cys	Val	
				325					330					335			
	ggc	cag	cac	gat	ttc	gat	ctc	ttg	agg	aga	ctc	gtt	ttg	cct	gac	ggg	1056

	Gly	Gln	His	Asp	Phe	Asp	Leu	Leu	Arg	Arg	Leu	Val	Leu	Pro	Asp	Gly	
				340					345					350			
	tcg	att	ttg	agg	tgt	gag	tac	tat	gct	ctc	cca	act	cgt	gac	cgt	ctc	1104
5	Ser	Ile	Leu	Arg	Cys	Glu	Tyr	Tyr	Ala	Leu	Pro	Thr	Arg	Asp	Arg	Leu	
			355					360					365				
	ttt	gaa	gac	cct	ctt	cat	gat	ggc	aaa	acc	atg	ctc	aag	att	tgg	aac	1152
	Phe	Glu	Asp	Pro	Leu	His	Asp	Gly	Lys	Thr	Met	Leu	Lys	Ile	Trp	Asn	
			370				375					380					
10	ttg	aac	aag	tac	act	gga	atc	atc	gga	gca	ttc	aac	tgt	caa	gga	gga	1200
	Leu	Asn	Lys	Tyr	Thr	Gly	Ile	Ile	Gly	Ala	Phe	Asn	Cys	Gln	Gly	Gly	
	385					390					395				400		
	gga	tgg	tgc	aga	gaa	act	cga	cgc	gac	caa	tgc	ttc	tcc	caa	tgc	gtt	1248
	Gly	Trp	Cys	Arg	Glu	Thr	Arg	Arg	Asp	Gln	Cys	Phe	Ser	Gln	Cys	Val	
				405					410					415			
15	aac	acg	tta	acc	gcc	aca	aca	aat	cct	aat	gac	gtt	gaa	tgg	aac	agt	1296
	Asn	Thr	Leu	Thr	Ala	Thr	Thr	Asn	Pro	Asn	Asp	Val	Glu	Trp	Asn	Ser	
				420					425					430			
	ggg	aac	aac	ccg	atc	tcc	att	gaa	aac	gtt	gaa	gag	ttt	gct	ttg	ttc	1344
20	Gly	Asn	Asn	Pro	Ile	Ser	Ile	Glu	Asn	Val	Glu	Glu	Phe	Ala	Leu	Phe	
			435					440					445				
	ttg	tct	caa	tcc	aag	aag	ctt	gtg	ttg	tcc	ggg	caa	aac	gat	gat	ctc	1392
	Leu	Ser	Gln	Ser	Lys	Lys	Leu	Val	Leu	Ser	Gly	Gln	Asn	Asp	Asp	Leu	
			450				455					460					
25	gag	atc	aca	tta	gag	ccc	ttc	aag	ttc	gag	ctc	atc	act	gtc	tca	cca	1440
	Glu	Ile	Thr	Leu	Glu	Pro	Phe	Lys	Phe	Glu	Leu	Ile	Thr	Val	Ser	Pro	
	465				470					475				480			
	gtt	gtc	acc	att	gag	ggc	agt	tcg	gtt	cag	ttt	gct	cca	atc	gga	ttg	1488
	Val	Val	Thr	Ile	Glu	Gly	Ser	Ser	Val	Gln	Phe	Ala	Pro	Ile	Gly	Leu	
				485					490				495				
30	gtt	aac	atg	ctt	aac	act	agc	ggt	gcg	att	cga	tcc	ttg	gtt	tat	cat	1536
	Val	Asn	Met	Leu	Asn	Thr	Ser	Gly	Ala	Ile	Arg	Ser	Leu	Val	Tyr	His	
				500					505					510			
	gag	gaa	tcc	gtt	gag	atc	ggt	gtt	cgt	ggt	gct	gga	gaa	ttc	agg	gtt	1584
35	Glu	Glu	Ser	Val	Glu	Ile	Gly	Val	Arg	Gly	Ala	Gly	Glu	Phe	Arg	Val	
			515					520					525				
	tat	gca	tcg	aag	aaa	cct	gtg	agc	tgc	aag	att	gat	ggt	gaa	gat	gtt	1632
	Tyr	Ala	Ser	Lys	Lys	Pro	Val	Ser	Cys	Lys	Ile	Asp	Gly	Glu	Asp	Val	
			530				535					540					
	gag	ttt	ggg	tac	gaa	gag	tca	atg	gtg	atg	gtt	caa	gtg	cct	tgg	tct	1680

Glu Phe Gly Tyr Glu Glu Ser Met Val Met Val Gln Val Pro Trp Ser  
545 550 555 560  
gca cca gag ggt ttg tct tct att aag tat ttg ttt tag agttatttaa 1729  
Ala Pro Glu Gly Leu Ser Ser Ile Lys Tyr Leu Phe  
5 565 570  
ggtgcttaat tgaaaaaaaa aaaaaaaaaa aaa 1762

<210> 9  
<211> 25  
10 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.  
15

<400> 9  
ccaatctgat catgcttgtg ccgaa 25

<210> 10  
20 <211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
25 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 10  
ggaacaaagt tatgcactat tatttaaggt 30

<210> 11  
30 <211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
35 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 11  
ctaccaaatt ccacaactta aagttca 27

<210> 12  
 <211> 32  
 <212> DNA  
 5 <213> Artificial Sequence  
  
 <220>  
 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.  
  
 10 <400> 12  
 ggaataataa gcttcacaca tactgtactc tc 32  
  
 <210> 13  
 <211> 30  
 15 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Designed oligonucleotide primer to obatin raffinose synthase gene.  
 20 <400> 13  
 atggctccaa gcttttagcaa ggaaaattcc 30  
  
 <210> 14  
 25 <211> 30  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 30 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.  
  
 <400> 14  
 tcaaaataag tactcaacag tggtaaaacc 30  
  
 35 <210> 15  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 15

5 ttggaagaga agacgccgcc gggaatcgtc 30

<210> 16

<211> 30

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 16

15 ttaagccccg gcgagagctc tggccggaca 30

<210> 17

<211> 30

20 <212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

25

<400> 17

accaatccaa aatctcatca aataatcgca 30

<210> 18

30 <211> 25

<212> DNA

<213> Artificial Sequence

<220>

35 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 18

aaataatagg ggcagtacaa attacaccac 30

<210> 19  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

5

<220>  
 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 19

10 atggctccac cgagcgtaat taaatccga 29

<210> 20

<211> 30

<212> DNA

15 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 20

20 ctaaaactca tacttaatag aagacaaacc 30

<210> 21

<211> 41

25 <212> DNA

<213> Artificial Sequence

<220>

30 <223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
 n is i.

<400> 21

cgatggatgg gnaanttnat ncancngan tggganatgt t 41

35 <210> 22

<211> 32

<212> DNA

<213> Artificial Sequence

Sequence



<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
n is i, r is a or g.

5

<400> 22

ggccacatnt tnacnarncc natngngcn aa

32

<210> 23

<211> 30

10

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

15

<400> 23

tggtactagg cgaaacaaga gtagctctga

30

<210> 24

20

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

25

<223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
n is i.

<400> 24

cgaggnggnt gncncnngg ntngtnatn atngangang gntggca

47

30

<210> 25

<211> 29

<212> DNA

<213> Artificial Sequence

35

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
n is i, y is t or c, r is a or g.

atytttrtcna cngcnarrtc ytccatngt

29

5

&lt;211&gt; 38

## <212> DNA

### <213> Artificial Sequence

 $\langle 220 \rangle$ 

10

<223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
n is i, y is t or c.

&lt;400&gt; 26

ggnacntant ggytncangg ntgncanatg gtncantg

38

15

&lt;210&gt; 27

&lt;211&gt; 32

## <212> DNA

### <213> Artificial Sequence

20

$\langle 220 \rangle$

<223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
n is 4, r is a or g.

25

&lt;400&gt; 27

ggccacatnt tnacnarncc natngngcn aa

32

&lt;210&gt; 28

&lt;211&gt; 26

30

## <212> DNA

### ⟨213⟩ Artificial Sequence

 $\langle 220 \rangle$ 

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

35

&lt;400&gt; 28

atctatttgt catgacgatg atccga

26

&lt;210&gt; 29

<211> 30  
<212> DNA  
<213> Artificial Sequence

5 <220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 29  
ggccctcatt cccatattgg gatgacctc 30

10 <210> 30  
<211> 30  
<212> DNA  
<213> Artificial Sequence

15 <220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 30

20 aagcatgcc aacatacaca tgctcaacag 30

<210> 31  
<211> 30  
25 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

30 <400> 31  
agaccgggg aaagctttgg ggttactact 30

<210> 32  
35 <211> 28  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 32

tggatgggaa actttataca ccctgact 28

5

<210> 33

<211> 28

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 33

15 gacatgttcc catctacaca cccttggtg 28

<210> 34

<211> 30

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

25 <400> 34

ccaatttatg ttagtgatgt tggtggcaag 30

<210> 35

<211> 26

30 <212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

35

<400> 35

tcgactccca gggtagaatt gtcacg 26

<210> 36

<211> 35  
<212> DNA  
<213> Artificial Sequence

5 <220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
n is i.

10 <400> 36  
cgattnaang tntggtggac nacncantgg gtngg 35

<210> 37  
<211> 38  
<212> DNA  
15 <213> Artificial Sequence

<220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene,  
n is i, r is a or g.

20 <400> 37  
cantgnacca tntgncancc ntgnarccan tangtncc 38

<210> 38  
25 <211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
30 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 38  
gttagggttc atatgaacac cttcaagctc 30

35 <210> 39  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 39

5 caacggcgag atcttgcac gtcaac 26

<210> 40

<211> 30

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 40

15 ggattcgaca caaacgcga cgtcatcgtc 30

<210> 41

<211> 27

20 <212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

25

<400> 41

ccacgtgcac caccgaact tatcgac 27

<210> 42

30 <211> 30

<212> DNA

<213> Artificial Sequence

<220>

35 <223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 42

aacatcgata ccatcggagt catgtccaat 30

<210> 43  
<211> 30  
<212> DNA  
<213> Artificial Sequence

5

<220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 43  
g t t a g g g t t c a t a t g a a c a c c t t c a a g c t c

30

<210> 44  
<211> 29  
<212> DNA  
<213> Artificial Sequence

15

<220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 44  
t c t a c g t c t g g c a c g c g c t t t g c g g c t a c

29

<210> 45  
<211> 31  
<212> DNA  
<213> Artificial Sequence

25

<220>  
<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

30

<400> 45  
g t t g a c g t c a t c c a c a t a t t g g a g a t g t t g t

31

<210> 46  
<211> 29  
<212> DNA  
<213> Artificial Sequence

35

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 46

gttatcgcta gcatggagca ctgtaatga 29

5

<210> 47

<211> 35

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

<400> 47

15 aacgagctca atccaaaatc tcatacaata atcgc 35

<210> 48

<211> 25

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to obtain raffinose synthase gene.

25 <400> 248

acaatagttg agggcggaag agtag 25

<210> 49

<211> 25

30 <212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide linker to obtain raffinose synthase gene.

35

<400> 49

gatcgagctc gtgtcggatc cagct 25

<210> 50



<211> 17  
 <212> DNA  
 <213> Artificial Sequence

5 <220>  
 <223> Designed oligonucleotide linker to obtain raffinose synthase gene.

<400> 50  
 ggatccgaca cgagctc 17

10

<210> 51  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

15

<220>  
 <223> Designed oligonucleotide primer to confirm direction of the inserted  
 raffinose synthase gene.

20 <400> 51  
 cctcctcgga ttccattgcc cagctatctg 30

<210> 52  
 <211> 30

25 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Designed oligonucleotide primer to confirm direction of the inserted  
 raffinose synthase gene.

30

<400> 52  
 ggattcgaca caaaccgcca cgtcacgctc 30

35 <210> 53  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

GenBank

<220>

<223> Designed oligonucleotide primer to confirm direction of the inserted  
raffinose synthase gene.

5

<400> 53

tctacgtctg gcacgcgctt tgcggctac

29

CCGCAGGCGCTT

What is claimed is:

1. A raffinose synthase gene which comprises a nucleotide sequence hybridizable with a nucleotide sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1,

(b) the nucleotide sequence represented by SEQ ID NO: 2,

10 (c) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 3,

(d) the nucleotide sequence represented by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4,

15 (e) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5,

(f) the nucleotide sequence represented by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6,

20 (g) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7, and

(h) the nucleotide sequence represented by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8,

25 under stringent conditions, and encoding a protein being capable of binding D-galactosyl group through  $\alpha$  (1 $\rightarrow$ 6) bond to the

hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule to form raffinose.

2. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1.

3. A raffinose synthase gene comprising the nucleotide sequence represented by SEQ ID NO: 2.

4. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 3.

5. A raffinose synthase gene comprising the nucleotide sequence represented by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4.

6. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5.

7. A raffinose synthase gene comprising the nucleotide sequence represented by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6.

8. A raffinose synthase gene comprising a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7.

9. A raffinose synthase gene comprising the

nucleotide sequence represented by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8.

10. A raffinose synthase gene comprising the nucleotide sequence represented by SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8.

11. A nucleic acid comprising a partial nucleotide sequence of the raffinose synthase gene of any one of claims 1 to 10.

12. A method for detecting a nucleic acid containing a raffinose synthase gene which comprises detecting said nucleic acid by hybridization using the labeled nucleic acid of claim 11 as a probe.

13. A method for amplifying a nucleic acid containing a raffinose synthase gene which comprises amplifying said nucleic acid by polymerase chain reaction (PCR) using the nucleic acid of claim 11 as a primer.

14. A method for obtaining a raffinose synthase gene which comprises the steps of:

detecting a nucleic acid containing said raffinose synthase gene by hybridization using the labeled nucleic acid of claim 11 as a probe, and

recovering the detected nucleic acid.

15. A method for obtaining a raffinose synthase gene which comprises the steps of:

amplifying a nucleic acid containing said raffinose

synthase gene by PCR using the nucleic acid of claim 11 as a primer, and

recovering the amplified nucleic acid.

5 16. A nucleic acid comprising a nucleic acid containing the raffinose synthase gene of any one of claims 1 to 10 ~~or the nucleic acid of claim 10~~ which is joined to a nucleic acid exhibiting promoter activity in a host cell.

April 16, 1999  
EW.

April 16, 1999  
K.O

17. A vector comprising the raffinose synthase gene of any one of claims 1 to 10.

10 18. A transformant, wherein the raffinose synthase gene of any one of claims 1 to 10 is introduced into a host cell.

19. A transformant, wherein the nucleic acid of claim 16 is introduced into a host cell.

15 20. A transformant, wherein the vector of claim 17 is introduced into a host cell.

21. The transformant of any one of claims 18 to 20, wherein the host is a microorganism.

22. The transformant of any one of claims 18 to 20, wherein the host is a plant.

20 23. A method for producing a raffinose synthase which comprises the steps of:

culturing or growing the transformant of any one of claims 18 to 22 to produce the raffinose synthase, and collecting the raffinose synthase.

25 24. A raffinose synthase comprising the amino acid

25. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 3.

27. A raffinose synthase comprising the amino acid sequence represented by SEQ ID NO: 7.

Abstract of the disclosure:

A raffinose synthase gene comprising a nucleotide sequence hybridizable with a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 1, (b) the nucleotide sequence represented by SEQ ID NO: 2, (c) a nucleotide sequence encoding the amino acid sequence of represented by SEQ ID NO: 3, (d) the nucleotide sequence represented by SEQ ID NO: 4 or by the 236th to 2584th nucleotides in the nucleotide sequence represented by SEQ ID NO: 4, (e) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 5, (f) the nucleotide sequence represented by SEQ ID NO: 6 or by the 134th to 2467th nucleotides in the nucleotide sequence represented by SEQ ID NO: 6, (g) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID NO: 7, and (h) the nucleotide sequence represented by SEQ ID NO: 8 or by the 1st to 1719th nucleotides in the nucleotide sequence represented by SEQ ID NO: 8, under stringent conditions, and encoding a protein being capable of binding D-galactosyl group through  $\alpha$  (1 $\rightarrow$ 6) bond to the hydroxyl group attached to the carbon atom at 6-position of the D-glucose residue in a sucrose molecule to form raffinose is disclosed.



# BIRCH, STEWART, KOLASCH & BIRCH, LLP

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

ATTORNEY DOCKET NO.  
20-4559P

PLEASE NOTE:  
YOU MUST  
COMPLETE THE  
FOLLOWING:

## COMBINED DECLARATION AND POWER OF ATTORNEY FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Insert Title: → RAFFINOSE SYNTHASE GENES AND THEIR USE

Fill in Appropriate  
Information —  
For Use  
Without  
Specification  
Attached:

the specification of which is attached hereto. If not attached hereto,

the specification was filed on \_\_\_\_\_ as  
United States Application Number \_\_\_\_\_;  
and amended on \_\_\_\_\_ (if applicable); and/or  
the specification was filed on \_\_\_\_\_ as PCT  
International Application Number \_\_\_\_\_; and was  
amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Insert Priority  
Information:  
(if appropriate)

Prior Foreign Application(s)			Priority Claimed	
<u>120550/1998</u> (Number)	<u>Japan</u> (Country)	<u>04/30/1998</u> (Month / Day / Year Filed)	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
<u>120551/1998</u> (Number)	<u>Japan</u> (Country)	<u>04/30/1998</u> (Month / Day / Year Filed)	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
<u>345590/1998</u> (Number)	<u>Japan</u> (Country)	<u>12/04/1998</u> (Month / Day / Year Filed)	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
<u>351246/1998</u> (Number)	<u>Japan</u> (Country)	<u>12/10/1998</u> (Month / Day / Year Filed)	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No

Insert Provisional  
Application(s):  
(if any)

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More than 12 Months (6 Months for Designs) Prior to the Filing Date of This Application:

Insert Requested  
Information:  
(if appropriate)

Country	Application Number	Date of Filing (Month / Day / Year)
_____	_____	_____
_____	_____	_____

I hereby claim the benefit under Title 35, United States Code, §120 of any United States and/or PCT application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States and/or PCT application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Insert Prior U.S.  
Application(s):  
(if any)

(Application Number)	(Filing Date)	(Status — patented, pending, abandoned)
_____	_____	_____
(Application Number)	(Filing Date)	(Status — patented, pending, abandoned)

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Raymond C. Stewart	(Reg. No. 21,066)	Terrell C. Birch	(Reg. No. 19,382)
Joseph A. Kolasch	(Reg. No. 22,463)	James M. Slattery	(Reg. No. 28,380)
Bernard L. Sweeney	(Reg. No. 24,448)	Michael K. Mutter	(Reg. No. 29,680)
Charles Gorenstein	(Reg. No. 29,271)	Gerald M. Murphy, Jr.	(Reg. No. 28,977)
Leonard R. Svensson	(Reg. No. 30,330)	Terry L. Clark	(Reg. No. 32,644)
Andrew D. Meikle	(Reg. No. 32,868)	Marc S. Weiner	(Reg. No. 32,181)
Joe McKinney Muncy	(Reg. No. 32,334)	C. Joseph Faraci	(Reg. No. 32,350)
Donald J. Daley	(Reg. No. 34,313)	John W. Bailey	(Reg. No. 32,881)
John A. Castellano	(Reg. No. 35,094)		

Send Correspondence to: **BIRCH, STEWART, KOLASCH & BIRCH, LLP**  
P.O. Box 747 • Falls Church, Virginia 22040-0747  
Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

**PLEASE NOTE:  
YOU MUST  
COMPLETE  
THE  
FOLLOWING:**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First or  
Sole Inventor:  
Insert Name of  
Inventor →  
Insert Date This  
Document is Signed

Insert Residence  
Insert Citizenship →

Insert Post Office  
Address →

Full Name of Second  
Inventor, if any:  
see above

Full Name of Third  
Inventor, if any  
see above

Full Name of Fourth  
Inventor, if any  
see above

Full Name of Fifth  
Inventor, if any  
see above

<b>GIVEN NAME</b>	<b>FAMILY NAME</b>	<b>INVENTOR'S SIGNATURE</b>	<b>DATE*</b>
Eijiro	WATANABE	<i>E. Watanabe</i>	April 16, 1999
Residence (City, State & Country)			<b>CITIZENSHIP</b>
Takarazuka-shi, Hyogo-ken, Japan			Japan
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
32-12-403, Fukui-cho, Takarazuka-shi, Hyogo-ken, Japan			
<b>GIVEN NAME</b>	<b>FAMILY NAME</b>	<b>INVENTOR'S SIGNATURE</b>	<b>DATE*</b>
Kenji	OEDA	<i>Kenji Oeda</i>	April 16, 1999
Residence (City, State & Country)			<b>CITIZENSHIP</b>
Kyoto-shi, Kyoto-fu, Japan			Japan
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
53-62-303, Shimogamoizumikawa-cho, Sakyo-ku, Kyoto-shi, Kyoto-fu, Japan			
<b>GIVEN NAME</b>	<b>FAMILY NAME</b>	<b>INVENTOR'S SIGNATURE</b>	<b>DATE*</b>
Residence (City, State & Country)			<b>CITIZENSHIP</b>
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
<b>GIVEN NAME</b>	<b>FAMILY NAME</b>	<b>INVENTOR'S SIGNATURE</b>	<b>DATE*</b>
Residence (City, State & Country)			<b>CITIZENSHIP</b>
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
<b>GIVEN NAME</b>	<b>FAMILY NAME</b>	<b>INVENTOR'S SIGNATURE</b>	<b>DATE*</b>
Residence (City, State & Country)			<b>CITIZENSHIP</b>
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			